

Store at
-20C
#65127**WHSC1 (D4Z8Q) Rabbit mAb**

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP, C&R	H Mk	Endogenous	66, 152	Rabbit IgG	#O96028	7468

Product Usage Information

The CUT&RUN dilution was determined using CUT&RUN Assay Kit #86652.

Application	Dilution
Western Blotting	1:1000
Immunoprecipitation	1:200
CUT&RUN	1:25

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

WHSC1 (D4Z8Z) Rabbit mAb recognizes endogenous levels of total WHSC1 protein. Based on antigen sequence, this antibody should detect WHSC1 Isoform II (UniProt O96028-1) and REIIBP (UniProt O96028-4). This antibody does not detect Isoform I (UniProt O96028-3).

Source / Purification

Monoclonal antibody is produced by immunizing animals with recombinant protein specific to the carboxy terminus of human WHSC1 protein.

Background

Wolf-Hirschhorn syndrome candidate protein 1 (WHSC1), also known as Nuclear SET domain-containing protein 2 (NSD2) and Multiple myeloma SET domain-containing protein (MMSET), is a histone methyltransferase protein that catalyzes the di-methylation of histone H3 Lys36, a mark that is associated with transcriptionally active regions of the genome (1,2). WHSC1 haploinsufficiency is implicated in the developmental disorder known as Wolf-Hirschhorn syndrome (WHS), which is characterized by growth and mental retardation, and congenital heart defects (3). In addition, WHSC1 is overexpressed and associated with poor prognosis in a large variety of human cancers, including neuroblastoma, and ovarian, hepatocellular, endometrial, and colorectal carcinoma (4-8). In multiple myeloma, 15-20% of all patients show overexpression of WHSC1 due to a t(4;14)(p16.3;q32) chromosomal translocation that places the *WHSC1* gene upstream of a strong enhancer (2). The *WHSC1* gene encodes for three major protein isoforms. Isoform II is the full-length protein containing 1365 amino acids. Isoform I, which is generated by alternative splicing, is identical to the N-terminal 647 amino acids of isoform II. A third isoform, known as REIIBP, is generated from a transcript initiating in intron 9 of the WHSC1 locus and is identical to the C-terminal 584 amino acids of isoform II.

Background References

1. Vougiouklakis, T. et al. (2015) *Epigenomics* 7, 863-74.
2. Dolloff, N.G. and Talamo, G. (2013) *Adv Exp Med Biol* 779, 197-221.
3. Stec, I. et al. (1998) *Hum Mol Genet* 7, 1071-82.
4. Hudlebusch, H.R. et al. (2011) *Clin Cancer Res* 17, 2919-33.
5. Yang, S. et al. (2013) *Biomarkers* 18, 257-63.
6. Zhou, P. et al. (2013) *Pathol Oncol Res* 19, 303-9.
7. Xiao, M. et al. (2013) *J Surg Oncol* 107, 428-32.
8. Hudlebusch, H.R. et al. (2011) *Cancer Res* 71, 4226-35.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation **C&R:** CUT&RUN

Cross-Reactivity Key

H: Human **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.