

#6604 Store at -20°C

SignalSilence® Atg7 siRNA I



✓ 10µM in 300 µl (100 transfections)

Orders ■ 877-616-CELL (2355) orders@cellsignal.com
Support ■ 877-678-TECH (8324) info@cellsignal.com
Web ■ www.cellsignal.com

rev. 02/16/16

For Research Use Only. Not For Use In Diagnostic Procedures.


Species Cross-Reactivity: H

Description: SignalSilence® Atg7 siRNA I from Cell Signaling Technology (CST) allows the researcher to specifically inhibit Atg7 expression using RNA interference, a method whereby gene expression can be selectively silenced through the delivery of double stranded RNA molecules into the cell. All SignalSilence® siRNA products from CST are rigorously tested in-house and have been shown to reduce target protein expression by western analysis.

Background: Autophagy is a catabolic process for the autophagosomal-lysosomal degradation of bulk cytoplasmic contents (1,2). Autophagy is generally activated by conditions of nutrient deprivation but has also been associated with a number of physiological processes including development, differentiation, neurodegeneration, infection and cancer (3). The molecular machinery of autophagy was largely discovered in yeast and referred to as **autophagy**-related (Atg) genes. Formation of the autophagosome involves an ubiquitin-like conjugation system in which Atg12 is covalently bound to Atg5 and targeted to autophagosome vesicles (4-6). This conjugation reaction is mediated by the ubiquitin-E1-like enzyme Atg7 and the E2-like enzyme Atg10 (7,8).

Directions for Use: CST recommends transfection with 100 nM Atg7 siRNA I 48 to 72 hours prior to cell lysis. For transfection procedure, follow protocol provided by the transfection reagent manufacturer. Please feel free to contact CST with any questions on use.

Quality Control: Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid phase extraction. The annealed RNA duplex is further analyzed by mass spectrometry to verify the exact composition of the duplex. Each lot is compared to the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency.



Western blot analysis of extracts from HeLa cells, transfected with 100 nM SignalSilence® Control siRNA (Unconjugated) #6568 (-) or SignalSilence® Atg7 siRNA I (+), using Atg7 Antibody #2631 (upper) or β-Tubulin (9F3) Rabbit mAb #2128 (lower). The Atg7 Antibody confirms silencing of Atg7 expression, while the β-Tubulin (9F3) Rabbit mAb is used as a loading control.

Entrez-Gene ID #10533
Swiss-Prot Acc. #O95352

Storage: Atg7 siRNA I is supplied in RNase-free water. Aliquot and store at -20°C.

Please visit www.cellsignal.com for a complete listing of recommended companion products.

Background References:

- (1) Reggiori, F. and Klionsky, D.J. (2002) *Eukaryot Cell* 1, 11-21.
- (2) Codogno, P. and Meijer, A.J. (2005) *Cell Death Differ* 12 Suppl 2, 1509-18.
- (3) Levine, B. and Yuan, J. (2005) *J Clin Invest* 115, 2679-88.
- (4) Mizushima, N. et al. (1998) *J Biol Chem* 273, 33889-92.
- (5) Mizushima, N. et al. (1998) *Nature* 395, 395-8.
- (6) Suzuki, K. et al. (2001) *EMBO J* 20, 5971-81.
- (7) Tanida, I. et al. (1999) *Mol Biol Cell* 10, 1367-79.
- (8) Shintani, T. et al. (1999) *EMBO J* 18, 5234-41.