

HCFC1 Antibody (Amino-terminal Antigen)

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP	Reactivity: H M R	Sensitivity: Endogenous	MW (kDa): 120, 130, 135, 160, 260	Source/Isotype: Rabbit	UniProt ID: #P51610	Entrez-Gene Id: 3054
-------------------------------	-----------------------------	-----------------------------------	--	----------------------------------	-------------------------------	--------------------------------

Product Usage Information	Application Western Blotting Immunoprecipitation	Dilution 1:1000 1:50
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.	
Specificity/Sensitivity	HCFC1 Antibody (Amino-terminal Antigen) recognizes endogenous levels of total HCFC1 protein. This antibody also recognizes amino-terminal fragments (HCFC1-N) resulting from O-GlcNAc transferase (OGT) cleavage.	
Source / Purification	Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Pro484 of human HCFC1 protein. Antibodies are purified by protein A and peptide affinity chromatography.	
Background	Host cell factor C1 (HCFC1) was first identified as the host cell factor for human herpes simplex virus infection. HCFC1 and the viral protein VP16 belong to a multi-protein complex that promotes transcription of viral immediate early genes (1). The relatively large HCFC1 protein contains 6 centrally located 26 amino acid repeats that can be O-GlcNAcylated and subjected to O-linked beta-N-acetylglucosamine transferase (OGT) cleavage (2-4). The resulting amino-terminal (HCFC1-N) and carboxy-terminal (HCFC1-C) fragments are non-covalently associated and play important roles in cell cycle regulation. The HCFC1-N peptide facilitates progression through the G1 phase of the cell cycle while HCFC1-C enables proper mitosis and cytokinesis during the M phase (5-7). As HCFC1 plays an important role in neurodevelopment, mutations in the corresponding gene are associated with neurodevelopmental disorders (e.g., intellectual disability) in humans (8).	
Background References	<ol style="list-style-type: none"> 1. Vogel, J.L. and Kristie, T.M. (2013) <i>Viruses</i> 5, 1272-91. 2. Daou, S. et al. (2011) <i>Proc Natl Acad Sci U S A</i> 108, 2747-52. 3. Capotosti, F. et al. (2011) <i>Cell</i> 144, 376-88. 4. Lazarus, M.B. et al. (2013) <i>Science</i> 342, 1235-9. 5. Julien, E. and Herr, W. (2003) <i>EMBO J</i> 22, 2360-9. 6. Julien, E. and Herr, W. (2004) <i>Mol Cell</i> 14, 713-25. 7. Zargar, Z. and Tyagi, S. (2012) <i>Transcription</i> 3, 187-92. 8. Jolly, L.A. et al. (2015) <i>Hum Mol Genet</i> 24, 3335-47. 	
Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).	
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.	
Applications Key	W: Western Blotting IP: Immunoprecipitation	
Cross-Reactivity Key	H: Human M: Mouse R: Rat	
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.	
Limited Uses	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's	

terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.