

**RALY Antibody**

**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W	H Mk	Endogenous	38	Rabbit	#Q9UKM9	22913

**Product Usage Information****Application**

Western Blotting

**Dilution**

1:1000

**Storage**

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

**Specificity/Sensitivity**

RALY Antibody recognizes endogenous levels of total RALY protein.

**Source / Purification**

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Pro163 of human RALY protein. Antibodies are purified by protein A and peptide affinity chromatography.

**Background**

RALY is a member of the large family of heterogeneous nuclear ribonucleoproteins (hnRNPs). RALY was initially discovered in lethal yellow mice, a condition where heterozygotes display an all yellow coat, obesity, diabetes, and tumors. In this condition, mRNA to the 5' untranslated region of RALY fuses to the agouti transcript (1,2). RALY binds to U-rich elements in coding and non-coding mRNAs undergoing translation (3). RALY has been implicated in certain cancer types, as it can control transcriptional regulation and splicing of E2F1 and PRMT1 mRNAs (4,5).

**Background References**

1. Michaud, E.J. et al. (1994) *Proc Natl Acad Sci U S A* 91, 2562-6.
2. Michaud, E.J. et al. (1993) *Genes Dev* 7, 1203-13.
3. Rossi, A. et al. (2017) *Nucleic Acids Res* 45, 6775-6792.
4. Bondy-Chorney, E. et al. (2017) *Int J Biochem Cell Biol* 91, 124-135.
5. Cornella, N. et al. (2017) *J Biol Chem* 292, 19674-19692.

**Species Reactivity**

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

**Western Blot Buffer**

**IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

**Applications Key**

**W:** Western Blotting

**Cross-Reactivity Key**

**H:** Human **Mk:** Monkey

**Trademarks and Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

**Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not

sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.