



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at +4C
#7687

E-Cadherin (24E10) Rabbit mAb (Alexa Fluor® 594 Conjugate)

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: IF-F, IF-IC	Reactivity: H M	Sensitivity: Endogenous	Source/Isotype: Rabbit IgG	UniProt ID: #P12830	Entrez-Gene Id: 999
-------------------------------------	---------------------------	-----------------------------------	--------------------------------------	-------------------------------	-------------------------------

Product Usage Information	Application Immunofluorescence (Frozen) Immunofluorescence (Immunocytochemistry)	Dilution 1:50 1:50
Storage	Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2 mg/ml BSA. Store at 4°C. Do not aliquot the antibody. Protect from light. Do not freeze.	
Specificity/Sensitivity	E-Cadherin (24E10) Rabbit mAb (Alexa Fluor® 594 Conjugate) recognizes endogenous levels of total E-cadherin protein. The antibody does not cross-react with related family members, such as N-cadherin.	
Species predicted to react based on 100% sequence homology	Bovine, Dog, Pig	
Source / Purification	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to the sequence surrounding Pro780 of human E-cadherin protein.	
Description	This Cell Signaling Technology antibody is conjugated to Alexa Fluor® 594 fluorescent dye and tested in-house for immunofluorescent analysis in human cells. This antibody is expected to exhibit the same species cross-reactivity as the unconjugated E-Cadherin (24E10) Rabbit mAb #3195.	
Background	<p>Cadherins are a superfamily of transmembrane glycoproteins that contain cadherin repeats of approximately 100 residues in their extracellular domain. Cadherins mediate calcium-dependent cell-cell adhesion and play critical roles in normal tissue development (1). The classic cadherin subfamily includes N-, P-, R-, B-, and E-cadherins, as well as about ten other members that are found in adherens junctions, a cellular structure near the apical surface of polarized epithelial cells. The cytoplasmic domain of classical cadherins interacts with β-catenin, γ-catenin (also called plakoglobin), and p120 catenin. β-catenin and γ-catenin associate with α-catenin, which links the cadherin-catenin complex to the actin cytoskeleton (1,2). While β- and γ-catenin play structural roles in the junctional complex, p120 regulates cadherin adhesive activity and trafficking (1-4). Investigators consider E-cadherin an active suppressor of invasion and growth of many epithelial cancers (1-3). Research studies indicate that cancer cells have upregulated N-cadherin in addition to loss of E-cadherin. This change in cadherin expression is called the "cadherin switch." N-cadherin cooperates with the FGF receptor, leading to overexpression of MMP-9 and cellular invasion (3). Research studies have shown that in endothelial cells, VE-cadherin signaling, expression, and localization correlate with vascular permeability and tumor angiogenesis (5,6). Investigators have also demonstrated that expression of P-cadherin, which is normally present in epithelial cells, is also altered in ovarian and other human cancers (7,8).</p>	
Background References	<ol style="list-style-type: none"> 1. Wheelock, M.J. and Johnson, K.R. (2003) <i>Annu Rev Cell Dev Biol</i> 19, 207-35. 2. Christofori, G. (2003) <i>EMBO J</i> 22, 2318-23. 3. Hazan, R.B. et al. (2004) <i>Ann N Y Acad Sci</i> 1014, 155-63. 4. Bryant, D.M. and Stow, J.L. (2004) <i>Trends Cell Biol</i> 14, 427-34. 5. Rabascio, C. et al. (2004) <i>Cancer Res</i> 64, 4373-7. 6. Yamaoka-Tojo, M. et al. (2006) <i>Arterioscler Thromb Vasc Biol</i> 26, 1991-7. 7. Patel, I.S. et al. (2003) <i>Int J Cancer</i> 106, 172-7. 8. Sanders, D.S. et al. (2000) <i>J Pathol</i> 190, 526-30. 	
Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).	
Applications Key	IF-F: Immunofluorescence (Frozen) IF-IC: Immunofluorescence (Immunocytochemistry)	
Cross-Reactivity Key	H: Human M: Mouse	

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

This product is provided under an intellectual property license from Life Technologies Corporation. The transfer of this product is conditioned on the buyer using the purchased product solely in research conducted by the buyer, excluding contract research or any fee for service research, and the buyer must not (1) use this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; or (c) manufacturing or quality assurance or quality control, and/or (2) sell or transfer this product or its components for resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.