

## Myelin Basic Protein (D8X4Q) XP® Rabbit



877-616-CELL (2355) Orders:

orders@cellsignal.com

Support: 877-678-TECH (8324)

info@cellsignal.com cellsignal.com Web:

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

## For Research Use Only. Not for Use in Diagnostic Procedures.

<b>Applications:</b> W, W-S, IHC-P, IF-F, IF-IC	Reactivity: H M R	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 12-18	<b>Source/Isotype:</b> Rabbit IgG	UniProt ID: #P02686	Entrez-Gene Id: 4155
Product Usage Information		Application Western Blotting			<b>Dilution</b> 1:1000	
		Simple Western™	to (Daraffin)			:50 - 1:250
		Immunohistochemis Immunofluorescence	•			:1200 :50
		Immunofluorescence	` '	nistry)		:50
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.				
		For a carrier free (BSA and azide free) version of this product see product #62596.				
Specificity/Sensitivity		Myelin Basic Protein (D8X4Q) XP <sup>®</sup> Rabbit mAb recognizes endogenous levels of total myelin basic protein.				
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ala185 of human myelin basic protein.				
Background		Myelin basic protein (MBP) is an abundant central nervous system (CNS) myelin membrane protein that plays an important role in nerve myelination. Myelin sheaths are multi-layered membranes derived from oligodendrocytes that increase the conduction velocity of axonal impulses. MBP helps to adhere the cytoplasmic leaflets of adjacent oligodendrocyte membranes to one another. Several splice variants of MBP are expressed in cells and the protein is modified following translation (i.e., phosphorylation, methylation), which suggests additional membrane adhesion functions (1).				
Background References		1. Harauz, G. and Bog	ງgs, J.M. (2013) <i>J Neເ</i>	urochem 125, 334-61.		
Species Reactiv	rity	Species reactivity is d	etermined by testin	g in at least one approve	ed application (e.g.,	western blot).
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.				
Applications Key		<b>W:</b> Western Blotting <b>W-S:</b> Simple Western™ <b>IHC-P:</b> Immunohistochemistry (Paraffin) <b>IF-F:</b> Immunofluorescence (Frozen) <b>IF-IC:</b> Immunofluorescence (Immunocytochemistry)				
Cross-Reactivit	y Key	H: Human M: Mouse	R: Rat			
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.				
		SignalStain is a registered trademark of Cell Signaling Technology, Inc.				
		XP is a registered trac	demark of Cell Signa	aling Technology, Inc.		
		All other trademarks more information.	are the property of	their respective owners.	Visit cellsignal.com	/trademarks for
Limited Uses		Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.				
				se Only or a similar labe		

purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com For Research Use Only. Not for Use in Diagnostic Procedures.