

#8144 Store at -20C

PDLIM2 Antibody



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H M	Sensitivity: Endogenous	MW (kDa): 38	Source/Isotype: Rabbit	UniProt ID: #Q96JY6	Entrez-Gene ID: 64236
Product Usage Information	Application Western Blotting			Dilution 1:1000		
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at - 20°C. Do not aliquot the antibody.					
Specificity/Sensitivity	PDLIM2 Antibody recognizes endogenous levels of total PDLIM2 protein.					
Source / Purification	Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Gly255 of human PDLIM2 protein. Antibodies are purified by protein A and peptide affinity chromatography.					
Background	PDLIM2, also known as Mystique, contains an amino-terminal PDZ domain and a carboxy-terminal LIM domain. PDLIM2 was originally found to be associated with cytoskeletal proteins in epithelial cells to promote cell attachment and migration (1,2). Subsequent studies have shown that PDLIM2 can also inhibit NF-κB activity by acting as a nuclear ubiquitin E3 ligase for p65 (3). PDLIM2 is suppressed in cancer cell lines by DNA methylation (4,5). Expression of PDLIM2 can inhibit anchorage-independent growth and tumor formation.					
Background References	1. Torrado, M. et al. (2004) <i>Invest Ophthalmol Vis Sci</i> 45, 3955-63. 2. Loughran, G. et al. (2005) <i>Mol Biol Cell</i> 16, 1811-22. 3. Tanaka, T. et al. (2007) <i>Nat Immunol</i> 8, 584-91. 4. Qu, Z. et al. (2010) <i>Cancer Res</i> 70, 1766-72. 5. Qu, Z. et al. (2010) <i>J Biol Chem</i> 285, 11786-92.					

Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.
Applications Key	W: Western Blotting
Cross-Reactivity Key	H: Human M: Mouse
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses	<p>Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.</p> <p>Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise</p>
---------------------	---

attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com
For Research Use Only. Not for Use in Diagnostic Procedures.