e at -20C	Met (D′	IC2) XP [®] Ra	bbit mAb				
Store						Orders:	877-616-CELL (2355) orders@cellsignal.com
						Support:	877-678-TECH (8324)
#8198						Web:	info@cellsignal.com cellsignal.com
#8					3 Trask Lane D	anvers Massao	chusetts 01923 USA
For Research Use Only. Not for Use in Diagnostic Procedures.							
	lications: -S, IP, IHC-	Reactivity: H	Sensitivity: Endogenous	MW (kDa): 140, 170	Source/Isotype: Rabbit IgG	UniProt ID: #P08581	Entrez-Gene Id: 4233

W, W-S, IP, IHC- H Bond, IHC-P, IF-IC, FC-FP	Endogenous	140, 170	Rabbit IgG	#P08581	4233		
Product Usage Information	Application Western Blotting Simple Western [™] Immunoprecipitation			Dilution 1:1000 1:50 - 1:2 1:50	250		
	IHC Leica Bond Immunohistochemist Immunofluorescence Flow Cytometry (Fixed	(Immunocytochem	istry)	1:150 - 1 1:150 - 1 1:1500 - 1:200 - 1	:600 1:3000		
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.						
	For a carrier-free (BSA	and azide free) ver	sion of this product see	product #83733.			
Specificity/Sensitivity	Met (D1C2) XP [®] Rabbi	t mAb recognizes e	ndogenous levels of tot	al Met protein.			
Source / Purification	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding residues near the carboxy terminus of human Met protein.				responding to		
Background	Met, a high affinity tyrosine kinase receptor for hepatocyte growth factor factor) is a disulfide-linked heterodimer made of 45 kDa α - and 145 kDa and the amino-terminal region of the β -subunit form the extracellular de chain spans the plasma membrane and contains a cytoplasmic region we Interaction of Met with HGF results in autophosphorylation at multiple to downstream signaling components, including Gab1, c-Cbl, and PI3 kinase events are important for all of the biological functions involving Met kinaphosphate at cytoplasmic Tyr1003 is essential for Met protein ubiquitina Phosphorylation at Tyr1234/1235 in the Met kinase domain is critical for Phosphorylation at Tyr1349 in the Met cytoplasmic domain provides a d Research studies have shown that altered Met levels and/or tyrosine kin several types of tumors, including renal, colon, and breast. Thus, investigi is an attractive potential cancer therapeutic and diagnostic target (6,7).			45 kDa β-subunits (1, ilular domain. The rer egion with tyrosines, whic I3 kinase (3). These fu Met kinase activity. Th iquitination and degr tical for kinase activat des a direct binding s sine kinase activities a investigators have co	2). The α-subunit nainder of the β- nase activity. h recruit several ndamental e addition of a adation (4). ion. ite for Gab1 (5). ire found in		
Background References	1. Cooper, C.S. et al. (1984) <i>Nature</i> 311, 29-33. 2. Bottaro, D.P. et al. (1991) <i>Science</i> 251, 802-4. 3. Bardelli, A. et al. (1997) <i>Oncogene</i> 15, 3103-11. 4. Taher, T.E. et al. (2002) <i>J Immunol</i> 169, 3793-800. 5. Schaeper, U. et al. (2000) <i>J Cell Biol</i> 149, 1419-32. 6. Eder, J.P. et al. (2009) <i>Clin Cancer Res</i> 15, 2207-14. 7. Sattler, M. and Salgia, R. (2009) <i>Update Cancer Ther</i> 3, 109-118.						
Species Reactivity	Species reactivity is de	etermined by testin	g in at least one approv	ed application (e.g., w	estern blot).		
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.						
Applications Key	W: Western Blotting W-S: Simple Western™ IP: Immunoprecipitation IHC-Bond: IHC Leica Bond IHC-P: Immunohistochemistry (Paraffin) IF-IC: Immunofluorescence (Immunocytochemistry) FC-FP: Flow Cytometry (Fixed/Permeabilized)						

Cross-Reactivity Key	H: Human				
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. XP is a registered trademark of Cell Signaling Technology, Inc.				
	All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.				
Limited Uses	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.				
	Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purpose, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.				