

#8337  
Store at -20C

## Glycolysis Antibody Sampler Kit

**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com**Support:** 877-678-TECH (8324)**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

1 Kit (8 x 20 microliters)

**For Research Use Only. Not for Use in Diagnostic Procedures.**

Product Includes	Product #	Quantity	Mol. Wt	Isotype/Source
PKM2 (D78A4) XP <sup>®</sup> Rabbit mAb	4053	20 µl	60 kDa	Rabbit IgG
GAPDH (D16H11) XP <sup>®</sup> Rabbit mAb	5174	20 µl	37 kDa	Rabbit IgG
Pyruvate Dehydrogenase (C54G1) Rabbit mAb	3205	20 µl	43 kDa	Rabbit IgG
Hexokinase I (C35C4) Rabbit mAb	2024	20 µl	102 kDa	Rabbit IgG
Hexokinase II (C64G5) Rabbit mAb	2867	20 µl	102 kDa	Rabbit
LDHA (C4B5) Rabbit mAb	3582	20 µl	37 kDa	Rabbit IgG
PKM1/2 (C103A3) Rabbit mAb	3190	20 µl	60 kDa	Rabbit IgG
PFKP (D4B2) Rabbit mAb	8164	20 µl	80 kDa	Rabbit IgG
Anti-rabbit IgG, HRP-linked Antibody	7074	100 µl		Goat

Please visit [cellsignal.com](http://cellsignal.com) for individual component applications, species cross-reactivity, dilutions, protocols, and additional product information.

### Description

The Glycolysis Antibody Sampler Kit provides an economical means to investigate select enzymes involved in glycolysis. The kit contains enough primary antibody to perform two western blot experiments with each primary antibody.

### Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

### Background

Glycolysis is the metabolic process by which glucose is converted to pyruvate in a sequence of enzymatic steps. Hexokinase catalyzes the conversion of glucose to glucose-6-phosphate, the first step in glycolysis. Hexokinases I, II, and III are associated with the outer mitochondrial membrane and are critical for maintaining an elevated rate of aerobic glycolysis in cancer cells (Warburg effect) (1). Phosphofructokinase (PFK) catalyzes the phosphorylation of fructose-6-phosphate. Platelet-type phosphofructokinase (PFKP) is expressed in various cell types (2). Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) catalyzes the phosphorylation of glyceraldehyde-3-phosphate (3). Pyruvate kinase, a glycolytic enzyme, catalyzes the conversion of phosphoenolpyruvate to pyruvate. In mammals, the M1 isoform (PKM1) is expressed in most adult tissues. The M2 isoform (PKM2), an alternatively-spliced variant of M1, is expressed during embryonic development (4). Lactate dehydrogenase (LDH) catalyzes the interconversion of pyruvate and NADH to lactate and NAD<sup>+</sup> (5). LDHA expression is induced when the oxygen supply is too low for mitochondrial ATP production (6). The pyruvate dehydrogenase complex catalyzes the conversion of pyruvate and CoA into acetyl-CoA and CO<sub>2</sub> in the presence of NAD<sup>+</sup>. The reaction of oxidative decarboxylation of pyruvate serves as a critical link between glycolysis and the citric acid cycle and lipid metabolism (7).

### Background References

1. WARBURG, O. (1956) *Science* 123, 309-14.
2. Morrison, N. et al. (1992) *Hum Genet* 89, 105-6.
3. Barber, R.D. et al. (2005) *Physiol Genomics* 21, 389-95.
4. Christofk, H.R. et al. (2008) *Nature* 452, 230-3.
5. Semenza, G.L. et al. (1996) *J Biol Chem* 271, 32529-37.
6. Semenza, G.L. (2007) *Biochem J* 405, 1-9.
7. Strumilo, S. (2005) *Acta Biochim Pol* 52, 759-64.

### Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

U.S. Patent No. 7,429,487, foreign equivalents, and child patents deriving therefrom.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

### Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless

separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

**Orders: 877-616-CELL (2355) • [orders@cellsignal.com](mailto:orders@cellsignal.com) • Support: 877-678-TECH (8324) • [info@cellsignal.com](mailto:info@cellsignal.com) • Web: [cellsignal.com](http://cellsignal.com)**  
For Research Use Only. Not for Use in Diagnostic Procedures.