

HSPA8 (D12F2) Rabbit mAb

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W	H M R Mk B Pg	Endogenous	70-72	Rabbit IgG	#P11142	3312

Product Usage Information**Application**

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

HSPA8 (D12F2) Rabbit mAb recognizes endogenous levels of total HSPA8 protein. This antibody cross-reacts with HSPA2, HSPA1A, and HSPC70.

Species predicted to react based on 100% sequence homology

Hamster, Chicken, D. melanogaster, Xenopus, Zebrafish, Dog, Horse

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Lys25 of human HSPA8 protein.

Background

HSPA8, alternately known as HSC70 or HSP73, is a constitutively expressed member of the HSP70 superfamily (1). Although its primary role in cells appears to be that of a general chaperone for unfolded proteins, HSPA8 has also been identified as the uncoating ATPase responsible for removing clathrin from coated vesicles and may also play a role in stabilizing untranslated mRNAs (1-5). In addition to these "housekeeping" functions, HSPA8 may also have an important role in inducible cellular stress responses. For example, oxidative or thermal stress promotes the nuclear/nucleolar accumulation of HSPA8, where it forms a complex with the topoisomerase I complex and likely protects it from heat inactivation (6,7). HSPA8 is reportedly phosphorylated in response to DNA damage, but it remains unclear what effect, if any, this has on HSPA8 function (8-10). Numerous high throughput studies support this observation. For more information, please see the HSPA8 page in PhosphoSitePlus® at www.phosphosite.org.

Background References

1. Takayama, S. et al. (1999) *J Biol Chem* 274, 781-6.
2. Goldfarb, S.B. et al. (2006) *Proc Natl Acad Sci USA* 103, 5817-22.
3. Cheetham, M.E. et al. (1996) *Biochem J* 319 (Pt 1), 103-8.
4. Ma, Y. et al. (2002) *J Biol Chem* 277, 49267-74.
5. Jønson, L. et al. (2007) *Mol Cell Proteomics* 6, 798-811.
6. Shiota, M. et al. (2010) *Hybridoma (Larchmt)* 29, 453-6.
7. Ciavarra, R.P. et al. (1994) *Proc Natl Acad Sci USA* 91, 1751-5.
8. Rush, J. et al. (2005) *Nat Biotechnol* 23, 94-101.
9. Matsuoka, S. et al. (2007) *Science* 316, 1160-6.
10. Beausoleil, S.A. et al. (2004) *Proc Natl Acad Sci USA* 101, 12130-5.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat **Mk:** Monkey **B:** Bovine **Pg:** Pig

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's

terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com
For Research Use Only. Not for Use in Diagnostic Procedures.