



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

# BCL6 (D4I2V) XP<sup>®</sup> Rabbit mAb (PE Conjugate)

Store at +4C  
#84882

**For Research Use Only. Not for Use in Diagnostic Procedures.**

<b>Applications:</b> FC-FP	<b>Reactivity:</b> H	<b>Sensitivity:</b> Endogenous	<b>Source/Isotype:</b> Rabbit IgG	<b>UniProt ID:</b> #P41182	<b>Entrez-Gene Id:</b> 604
-------------------------------	-------------------------	-----------------------------------	--------------------------------------	-------------------------------	-------------------------------

## Product Usage Information

### Application

Flow Cytometry (Fixed/Permeabilized)

### Dilution

1:50

## Storage

Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2 mg/ml BSA. Store at 4°C. Do not aliquot the antibodies. Protect from light. Do not freeze.

## Specificity/Sensitivity

BCL6 (D4I2V) XP<sup>®</sup> Rabbit mAb (PE Conjugate) recognizes endogenous levels of total BCL6 protein. It is predicted to detect both human BCL6 isoform 1 and isoform 2.

## Source / Purification

Monoclonal antibody is produced by immunizing animals with a recombinant protein fragment spanning residues Asp210 through Thr409 of human BCL6 protein.

## Description

This Cell Signaling Technology antibody is conjugated to phycoerythrin (PE) and tested in-house for direct flow cytometry analysis in human cells. This antibody is expected to exhibit the same species cross-reactivity as the unconjugated BCL6 (D4I2V) XP<sup>®</sup> Rabbit mAb #14895.

## Background

Chromosomal translocations result in misregulation of the proto-oncogene *BCL6* in patients with B cell-derived non-Hodgkin's lymphoma (1). The *BCL6* gene is selectively expressed in mature B cells and encodes a nuclear phosphoprotein that belongs to the BTB/POZ zinc finger family of transcription factors (2,3). BCL6 protein can bind to target DNA sequences of Stat6 and, analogous to Stat6, modulate the expression of interleukin-4-induced genes (4). Furthermore, BCL6 restrains p53-dependent senescence, making BCL6-active tumors functionally p53-negative (5). The mitogen-activated protein kinases, Erk1 and Erk2, but not JNK, phosphorylate BCL6 at multiple sites. Phosphorylation of BCL6 at Ser333 and Ser343 results in degradation of BCL6 by the ubiquitin/proteasome pathway in B cells (6,7). In addition, BCL6 is acetylated and its transcriptional repressor function is inhibited by the transcriptional co-activator p300 (8).

## Background References

1. Ye, B.H. et al. (1993) *Science* 262, 747-50.
2. Onizuka, T. et al. (1995) *Blood* 86, 28-37.
3. Zollman, S. et al. (1994) *Proc Natl Acad Sci U S A* 91, 10717-21.
4. Harris, M.B. et al. (1999) *Mol Cell Biol* 19, 7264-75.
5. Shvarts, A. et al. (2002) *Genes Dev* 16, 681-6.
6. Moriyama, M. et al. (1997) *Oncogene* 14, 2465-74.
7. Niu, H. et al. (1998) *Genes Dev* 12, 1953-61.
8. Bereshchenko, O.R. et al. (2002) *Nat Genet* 32, 606-13.

## Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

## Applications Key

**FC-FP:** Flow Cytometry (Fixed/Permeabilized)

## Cross-Reactivity Key

**H:** Human

## Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

## Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no

force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.