

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

#8496 Store at -20C

Phospho-Phospholamban (Ser16/Thr17) Antibody

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: R	Sensitivity: Endogenous	MW (kDa): 6 (monomer); 12, 24 (oligomers)	Source/Isotype: Rabbit
---------------------------	-------------------------	-----------------------------------	---	----------------------------------

Product Usage Information

Application

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

Phospho-Phospholamban (Ser16/Thr17) Antibody recognizes endogenous levels of phospholamban protein only when phosphorylated at Ser16 and Thr17. This antibody does not detect mono- or non-phosphorylated phospholamban.

Species predicted to react based on 100% sequence homology

Human, Mouse, Bovine, Dog, Pig

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Ser16/Thr17 of human phospholamban protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

Phospholamban (PLN) was identified as a major phosphoprotein component of the sarcoplasmic reticulum (SR) (1). Its name, "lamban", is derived from the greek word "lambano" meaning "to receive", so named due to the fact that phospholamban is heavily phosphorylated on serine and threonine residues in response to cardiac stimulation (1). Although originally thought to be a single 20-25 kDa protein due to its electrophoretic mobility on SDS-PAGE, PLN is actually a 52 amino acid, 6 kDa, membrane-spanning protein capable of forming stable homooligomers, even in the presence of SDS (2). Despite very high expression in cardiac tissue, phospholamban is also expressed in skeletal and smooth muscle (3). Localization of PLN is limited to the SR, where it serves as a regulator of the sarco-endoplasmic reticulum calcium ATPase, SERCA (4). PLN binds directly to SERCA and effectively lowers its affinity for calcium, thus reducing calcium transport into the SR. Phosphorylation of PLN at Ser16 by Protein Kinase A or myotonic dystrophy protein kinase and/or phosphorylation at Thr17 by Ca²⁺/calmodulin-dependent protein kinase results in release of PLN from SERCA, relief of this inhibition, and increased calcium uptake by the SR (reviewed in 5,6). It has long been held that phosphorylation at Ser16 and Thr17 occurs sequentially, but increasing evidence suggests that phosphorylation, especially at Thr17, may be differentially regulated (reviewed in 7,8).

Rodent models of heart failure have shown that the expression level and degree of phosphorylation of PLN are critical in modulating calcium flux and contractility (reviewed in 9-11). Deletion or decreased expression of PLN promotes increased calcium flux and increased cardiac contractility, whereas overexpression of PLN results in sequestration of SERCA, decreased calcium flux, reduced contractility, and rescue of cardiac dysfunction and failure in mouse models of hypertension and cardiomyopathy (reviewed in 10). Distinct mutations in PLN have been detected in humans, resulting either in decreased or no expression of PLN protein (12,13) or binding defects between PLN, SERCA and/or regulatory proteins (14,15), both of which result in cardiac myopathy and heart failure. Interestingly, while the human phenotype of most PLN defects mimic those seen in rodent and vice versa, there are some instances where the type and severity of cardiac disease resulting from PLN mutations in rodent and human differ, making a consensus mechanism elusive.

Background References

- Kirchberber, M.A. et al. (1975) *Recent Adv Stud Cardiac Struct Metab* 5, 103-15.
- Zhan, Q.Q. et al. (1991) *J Biol Chem* 266, 21810-4.
- Fujii, J. et al. (1991) *J Biol Chem* 266, 11669-75.
- Tada, M. and Kirchberger, M.A. *Recent Adv Stud Cardiac Struct Metab* 11, 265-72.
- Traaseth, N.J. et al. (2008) *Biochemistry* 47, 3-13.
- Bhupathy, P. et al. (2007) *J Mol Cell Cardiol* 42, 903-11.
- Hagemann, D. and Xiao, R.P. (2002) *Trends Cardiovasc Med* 12, 51-6.
- Mattiazzi, A. et al. (2005) *Cardiovasc Res* 68, 366-75.

9. Chu, G. and Kranias, E.G. (2006) *Novartis Found Symp* 274, 156-71; discussion 172-5, 272-6.
 10. Schwinger, R.H. and Frank, K.F. (2003) *Sci STKE* 2003, pe15.
 11. MacLennan, D.H. and Kranias, E.G. (2003) *Nat Rev Mol Cell Biol* 4, 566-77.
 12. Haghghi, K. et al. (2008) *Hum Mutat* 29, 640-7.
 13. Haghghi, K. et al. (2003) *J Clin Invest* 111, 869-76.
 14. Schmitt, J.P. et al. (2003) *Science* 299, 1410-3.
 15. Haghghi, K. et al. (2006) *Proc Natl Acad Sci USA* 103, 1388-93.
-

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting

Cross-Reactivity Key

R: Rat

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.