



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C
#8502

PEN2 (D2G6) Rabbit mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP	Reactivity: H M R Mk	Sensitivity: Endogenous	MW (kDa): 13	Source/Isotype: Rabbit IgG	UniProt ID: #Q9NZ42	Entrez-Gene Id: 55851
-------------------------------	--------------------------------	-----------------------------------	------------------------	--------------------------------------	-------------------------------	---------------------------------

Product Usage Information

Application

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

PEN2 (D2G6) Rabbit mAb recognizes endogenous levels of total PEN2 protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the amino terminus of human PEN2 protein.

Background

Presenilin Enhancer 2 (PEN2) is a small integral membrane glycoprotein that contains two recognized transmembrane domains. Both the N- and C-terminal domains are oriented into the lumen of the endoplasmic reticulum (1). PEN2, along with Presenilin 1, Presenilin 2, Nicastrin, and APH-1 form the protein complex γ -secretase (2). The proteinase BACE catalyses the initial step in APP processing by cleaving and releasing soluble APP β (3). The remaining membrane bound APP is then cleaved by the γ -secretase complex, causing the release of amyloid β -peptide, the main constituent of amyloid plaques. These plaques are a hallmark of Alzheimer's disease pathology (2). In addition to APP, the γ -secretase complex cleaves several other proteins and necessary presenilin-dependent signaling cascades, including the Notch pathway (4). It was found that PEN2 is an important part of the γ -secretase complex, and knocking it down results in reduced amounts of the complex, resulting in a loss of γ -secretase activity (5).

Background References

1. Sala Frigerio, C. et al. (2005) *J Neurol* 252, 1033-6.
2. Hansson, C.A. et al. (2004) *J Biol Chem* 279, 51654-60.
3. Hunt, C.E. and Turner, A.J. (2009) *FEBS J* 276, 1845-59.
4. St George-Hyslop, P. and Schmitt-Ulms, G. (2010) *Nature* 467, 36-7.
5. Steiner, H. et al. (2002) *J Biol Chem* 277, 39062-5.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in

any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.