

Stargazin (D33G3) Rabbit mAb

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP	M R	Endogenous	40	Rabbit IgG	#Q9Y698	10369

Product Usage Information**Application**

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

Stargazin (D33G3) Rabbit mAb recognizes endogenous levels of total stargazin protein.

Species predicted to react based on 100% sequence homology

Human

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ala210 of human stargazin protein.

Background

Stargazin is a four-pass transmembrane protein related to VDCC (voltage dependent calcium channel) γ subunits and part of the TARP (transmembrane AMPA receptor regulatory protein) family of proteins. TARP proteins can form a complex with AMPA receptors (GluR1-4) and serve as integral auxiliary subunits (1-6).

Interactions between stargazin and AMPA receptors are implicated in regulation of receptor surface expression, synaptic clustering and recycling, as well as increased receptor responsiveness to glutamate (1,2,5,6). Stargazin may play a role in the molecular mechanism of AMPAR-mediated inflammatory pain by taking part in signaling pathways that relay pain in the spinal cord (5). Because the protein also modulates the pharmacology of AMPA receptors, it enhances the effects of AMPAR potentiators that have therapeutic potential for a number of mental and neurodegenerative diseases (6).

The carboxy terminus of the stargazin protein interacts with the PDZ domains of PSD95 and other membrane-associated guanylate kinase (MAGUK) family members, and together traffic AMPA receptors to the cell surface membrane, anchoring them to the postsynaptic site (1,7). Phosphorylation of stargazin by PKA on Thr321 inhibits this binding (3).

Background References

- Tomita, S. et al. (2005) *Nature* 435, 1052-1058.
- Priel, A. et al. (2005) *J. Neurosci.* 25, 2682-2686.
- Choi, J. et al. (2002) *J. Biol. Chem.* 277, 12359-12363.
- Osten, P. and Stern-Bach, Y. (2006) *Curr. Opin. Neurobiol.* 16, 275-280.
- Tao, F. et al. (2006) *J. Neurosci. Res.* 84, 867-873.
- Tomita, S. et al. (2006) *Proc. Natl. Acad. Sci. USA* 103, 10064-10067.
- Schnell, E. et al. (2002) *Proc. Natl. Acad. Sci. USA* 99, 13902-13907.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

M: Mouse **R:** Rat

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.