8533

PDGF Receptor α (D13C6) XP[®] Rabbit mAb (PE Conjugate)



Orders:	877-616-CELL (2355) orders@cellsignal.com
Support:	877-678-TECH (8324)
Web:	info@cellsignal.com cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: FC-FP, FC-L	Reactivity: H	Sensitivity: Endogenous	Source/Isotype: Rabbit IgG	UniProt ID: #P16234	Entrez-Gene Id: 5156	
Product Usage Information		Application Flow Cytometry (Fixed/P Flow Cytometry (Live)		ride and 2 mg/ml BC	Dilution 1:50 1:50	
Storage		Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2 mg/ml BSA. Store at 4°C. Do not aliquot the antibodies. Protect from light. Do not freeze.				
Specificity/Sensit	City/Sensitivity PDGF Receptor α (D13C6) XP [®] Rabbit mAb detects endogenous levels of PDGF receptor α protein.					
Source / Purifica	tion	Monoclonal antibody is produced by immunizing animals with a recombinant protein corresponding the PDGF receptor α extracellular domain.				
Description		This Cell Signaling Technology antibody is conjugated to phycoerythrin (PE) and tested in-house for direct flow cytometry analysis in human cells. The antibody is expected to exhibit the same species cross-reactivity as the unconjugated PDGF Receptor α (D13C6) XP [®] Rabbit mAb #5241.				
Background		Platelet derived growth factor (PDGF) family proteins exist as several disulphide-bonded, dimeric isoforms (PDGF AA, PDGF AB, PDGF BB, PDGF CC, and PDGF DD) that bind in a specific pattern to two closely related receptor tyrosine kinases, PDGF receptor α (PDGFRα) and PDGF receptor β (PDGFRβ). PDGFRα and PDGFRβ share 75% to 85% sequence homology between their two intracellular kinase domains, while the kinase insert and carboxy-terminal tail regions display a lower level (27% to 28%) of homology (1). PDGFRα homodimers bind all PDGF isoforms except those containing PDGF D. PDGFRβ homodimers bind PDGF BB and DD isoforms, as well as the PDGF AB heterodimer. The heteromeric PDGF receptor α/β binds PDGF B, C, and D homodimers, as well as the PDGF AB heterodimer (2). PDGFRα and PDGFRβ can each form heterodimers with EGFR, which is also activated by PDGF (3). Various cells differ in the total number of receptors present and in the receptor subunit composition, which may account for responsive differences among cell types to PDGF binding (4). Ligand binding induces receptor dimerization and autophosphorylation, followed by binding and activation of cytoplasmic SH2 domain-containing signal transduction molecules, such as GRB2, Src, GAP, P13 kinase, PLCγ, and NCK. A number of different signaling pathways are initiated by activated PDGF receptors and lead to control of cell growth, actin reorganization, migration, and differentiation (5). Tyr751 in the kinase-insert region of PDGFRβ (pTyr751-VaI-Pro-Met-Leu) inhibit the association of the carboxy-terminal SH2 domain of the p85 subunit of P13 kinase with PDGFRβ (7). Tyr740 is also required for PDGFRβ-mediated P13 kinase activation (8).				
Background Refe	erences	 Deuel, T.F. et al. (1988) <i>Biofactors</i> 1, 213-217. Bergsten, E. et al. (2001) <i>Nat. Cell Biol.</i> 3, 512-516. Betsholtz, C. et al. (2001) <i>Bioessays</i> 23, 494-507. Coughlin, S.R. et al. (1988) <i>Prog. Clin. Biol. Res.</i> 266, 39-45. Ostman, A. and Heldin, C.H. (2001) <i>Adv. Cancer Res.</i> 80, 1-38. Panayotou, G. et al. (1992) <i>EMBO J.</i> 11, 4261-4272. Ramalingam, K. et al. (1992) <i>EMBO J.</i> 11, 1373-1382. 				
Species Reactivit	у	Species reactivity is determined by testing in at least one approved application (e.g., western blot).				
Applications Key		FC-FP: Flow Cytometry (Fixed/Permeabilized) FC-L: Flow Cytometry (Live)				
Cross-Reactivity Key H: Human						
Trademarks and Patents Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.				Inc.		

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.