



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

CA2 Antibody

Store at -20C
#8612

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP	Reactivity: H	Sensitivity: Endogenous	MW (kDa): 28	Source/Isotype: Rabbit	UniProt ID: #P00918	Entrez-Gene Id: 760
-------------------------------	-------------------------	-----------------------------------	------------------------	----------------------------------	-------------------------------	-------------------------------

Product Usage Information

Application

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

CA2 Antibody recognizes endogenous levels of total CA2 protein.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Lys169 of human CA2 protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

Carbonic anhydrases (CA) are a family of ancient zinc metalloenzymes found in almost all living organisms. All CA can be divided into 3 distinct classes (α , β , and γ) that evolved independently and have no significant homology in sequence and overall folding. All functional CA catalyze the reversible hydration of CO_2 into HCO_3^- and H^+ and contain a zinc atom in the active sites essential for catalysis. There are many isoforms of CA in mammals and they all belong to the α class (1,2). CA2 is a cytosolic member of the α class. It is the most widely distributed isoform among the mammalian CAs (1). Defects in CA2 are associated with osteopetrosis and renal tubular acidosis (3-5). Elevated expression of CA2 is observed in patients with Alzheimer's disease and the developing brains of Down syndrome patients (6,7). CA2 is also overexpressed in Gastrointestinal Stromal Tumors (GISTs) and is considered a useful marker for diagnosis (8). Recently, CA2 was reported to facilitate transporter activity of the monocarboxylate transporter isoform 1 and 4 (MCT1/4) independent of its own catalytic activity (9,10)

Background References

- Smith, K.S. et al. (1999) *Proc Natl Acad Sci USA* 96, 15184-9.
- Tripp, B.C. et al. (2001) *J Biol Chem* 276, 48615-8.
- McMahon, C. et al. (2001) *Blood* 97, 1947-50.
- Borthwick, K.J. et al. (2003) *J Med Genet* 40, 115-21.
- Lai, L.W. et al. (1998) *J Clin Invest* 101, 1320-5.
- Palminiello, S. et al. (2008) *Brain Res* 1190, 193-205.
- Jang, B.G. et al. (2010) *J Alzheimers Dis* 21, 939-45.
- Parkkila, S. et al. (2010) *Mod Pathol* 23, 743-50.
- Becker, H.M. et al. (2005) *J Biol Chem* 280, 39882-9.
- Becker, H.M. et al. (2011) *Proc Natl Acad Sci USA* 108, 3071-6.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: Human

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.