

Store at
-20C
#86668**MFF Antibody**

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP	H M	Endogenous	25, 27, 30, 35	Rabbit	#Q9GZY8	56947

Product Usage Information**Application**

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:200

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

MFF Antibody recognizes endogenous levels of total MFF protein. Based upon sequence alignment, this antibody is predicted to react with isoforms 1-5 of human MFF protein.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Lys52 of human MFF protein, isoform 1. Antibodies are purified by protein A and peptide affinity chromatography.

Background

Mitochondrial fission factor (MFF) is a tail-anchored protein that resides within the outer mitochondrial membrane and is part of the mitochondrial fission complex. MFF participates in mitochondrial fission by serving as one of multiple receptors for the GTPase dynamin-related protein 1 (Drp1) (1-4). Research studies have also shown that MFF is a peroxisomal membrane protein and participates in peroxisome fission by serving as a receptor for another GTPase, dynamin-like protein 1 (5,6). Research studies have demonstrated that the ability of MFF to drive acute mitochondrial fission in response to mitochondrial stress is controlled by AMPK-dependent phosphorylation. AMPK directly phosphorylates MFF at two sites to allow for enhanced recruitment of Drp1 to the mitochondria (7). Multiple isoforms of MFF are expressed as a result of alternative splicing (8). One of the major phosphoacceptor sites of MFF (Ser172 of human isoform 1/Ser146 of human isoforms 2-5) lies within an AMPK phosphorylation motif that spans the boundary of differentially spliced exons of MFF isoforms, suggesting that MFF splice isoforms are phosphorylated by AMPK to varying degrees.

Background References

1. Liu, R. and Chan, D.C. (2015) *Mol Biol Cell* 26, 4466-77.
2. Shen, Q. et al. (2014) *Mol Biol Cell* 25, 145-59.
3. Losón, O.C. et al. (2013) *Mol Biol Cell* 24, 659-67.
4. Otera, H. et al. (2010) *J Cell Biol* 191, 1141-58.
5. Itoyama, A. et al. (2013) *Biol Open* 2, 998-1006.
6. Gandre-Babbe, S. and van der Bliek, A.M. (2008) *Mol Biol Cell* 19, 2402-12.
7. Toyama, E.Q. et al. (2016) *Science* 351, 275-81.
8. Ducommun, S. et al. (2015) *Cell Signal* 27, 978-88.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: Human **M:** Mouse

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless

separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com
For Research Use Only. Not for Use in Diagnostic Procedures.