

Store at +4C  
#8726**Rabbit (DA1E) mAb IgG XP<sup>®</sup> Isotype Control (Magnetic Bead Conjugate)**

**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

<b>Applications:</b>	<b>Sensitivity:</b>	<b>Source/Isotype:</b>				
IP	Endogenous	Rabbit IgG				
<b>Product Usage Information</b>	<p>Directions for Use: Add 10 µl of well-vortexed beads to 200 µl of cell lysate at 1 mg/ml in 1X Cell Lysis Buffer (10X) #9803. See protocol for more details.</p> <p>For bead washing and subsequent elution of immunocomplexes, the beads can be separated from solution using our 6-Tube Magnetic Separation Rack #7017. Place the tubes containing the beads in the Magnetic Separation Rack and wait 1 to 2 minutes for the solution to clear before carefully removing the supernatant. Remove the tubes from the Magnetic Separation Rack, add new solution and resuspend the beads by gently vortexing or rocking the tube.</p> <table border="1"> <thead> <tr> <th><b>Application</b></th> <th><b>Dilution</b></th> </tr> </thead> <tbody> <tr> <td>Immunoprecipitation</td> <td>1:20</td> </tr> </tbody> </table>		<b>Application</b>	<b>Dilution</b>	Immunoprecipitation	1:20
<b>Application</b>	<b>Dilution</b>					
Immunoprecipitation	1:20					
<b>Storage</b>	Supplied in PBS Buffer (pH 7.2), 0.1% Tween <sup>®</sup> 20. Store at 4°C. Do not aliquot the antibodies.					
<b>Specificity/Sensitivity</b>	Rabbit (DA1E) mAb IgG XP <sup>®</sup> Isotype Control (Magnetic Bead Conjugate) is not directed against any known antigen. It functions as an isotype control for rabbit IgG monoclonal antibodies.					
<b>Description</b>	This Cell Signaling Technology (CST) antibody is immobilized by the covalent reaction of hydrazinonicotinamide-modified antibody with formylbenzamide-modified magnetic bead. Rabbit (DA1E) mAb IgG XP <sup>®</sup> Isotype Control (Magnetic Bead Conjugate) is useful to determine non-specific immunoprecipitation complexes.					
<b>Background</b>	Isotype control antibodies are used to estimate the nonspecific binding of target primary antibodies due to F <sub>c</sub> receptor binding or other protein-protein interactions. An isotype control antibody should have the same immunoglobulin type and be used at the same concentration as the test antibody.					
<b>Species Reactivity</b>	Species reactivity is determined by testing in at least one approved application (e.g., western blot).					
<b>Applications Key</b>	IP: Immunoprecipitation					
<b>Trademarks and Patents</b>	<p>Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.</p> <p>XP is a registered trademark of Cell Signaling Technology, Inc.</p> <p>All other trademarks are the property of their respective owners. Visit <a href="http://cellsignal.com/trademarks">cellsignal.com/trademarks</a> for more information.</p>					
<b>Limited Uses</b>	<p>Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.</p> <p>Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the</p>					

purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.