

8763

A2B5 Mouse mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

| Applications: F IF-IC | Reactivity: | Sensitivity: Endogenous | Source/Isotype: Mouse IgM | |
|------------------------------|-------------|--|---|--------------------------------|
| Product Usage Information | | It is strongly recommended that these directions be followed for staining of live, unpermeabilized cells when using this antibody. Dilute primary antibody 1:100 in culture medium containing 0.1% sodium azide*. Aspirate cell culture medium, add diluted antibody and incubate cells for 30 minutes at 37°C. Aspirate antibody solution, rinse gently with pre-warmed PBS, and fix with methanol-free, 4% formaldehyde diluted in PBS for 15 minutes at 37°C. Rinse cells with PBS 3 times for 5 minutes each. For best results, detect with a fluorochrome-conjugated anti-mouse IgM secondary antibody (per manufacturer's recommendation, diluted in PBS). Prior to imaging, rinse samples in PBS 3 times for 5 minutes each and mount in an anti-fade reagent (ProLong® Gold Antifade Reagent #9071). | | |
| | | * sodium azide is included to prevent capping. | | |
| | | Application Immunofluorescence (Ir | nmunocytochemistry) | Dilution 1:100 |
| Storage | | Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody. | | |
| Specificity/Sensitivity | ty | A2B5 Mouse mAb recognizes endogenous levels of A2B5 antigen. A2B5 Mouse mAb has been validated for use with the recommended staining protocol. Fixation of cells prior to staining will lead to artifacts. Based on literature publications, this antibody is predicted to cross-react with chicken, rat, bovine, and human samples. | | |
| Source / Purification | 1 | Monoclonal antibody is produced by immunizing animals with embryonic chicken retinal cells. | | |
| Background | | A2B5 Mouse mAb recognizes a cell surface ganglioside epitope that has been utilized as a marker for identification of various cell types. A2B5 Mouse mAb has been used to mark specific cell populations such as neuroendocrine cells, thymic epithelial cells (1), and glial precursors that give rise to type II astrocytes and oligodendrocytes (2-4). | | |
| Background References | | Botham, C.A. et al. (2001) Cell Tissue Res 303, 381-9. Eisenbarth, G.S. et al. (1979) Proc Natl Acad Sci U S A 76, 4913-7. Sontheimer, H. et al. (1990) Proc Natl Acad Sci U S A 87, 9833-7. Bottenstein, J.E. et al. (1988) J Neurosci Res 20, 291-303. | | |
| Species Peactivity | | Species reactivity is dete | rmined by tecting in at least one approved an | polication (e.g. western blot) |

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Applications Key

IF-IC: Immunofluorescence (Immunocytochemistry)

Cross-Reactivity Key

R: Rat

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in

any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.