CD8α (2.43) Rat mAb (PE-Cy7® Conjugate)



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: FC-L	Reactivity: M	Sensitivity: Endogenous	Source/Isotype: Rat IgG2b	UniProt ID: #P01731	Entrez-Gene Id: 12525
Product Usage		For optimal flow cytometry results, we recommend 0.25 μg of antibody per test.			
Information T		Application Dilution Flow Cytometry (Live) 1:80			
Storage		Supplied in 10 mM NaH $_2$ PO $_4$, 150 mM NaCl, 0.09% NaN $_3$, 0.1% gelatin, pH 7.2. This product is stable for 6 months when stored at 4 $^\circ$ C. Do not aliquot the antibody. Protect from light. Do not freeze.			
Specificity/Sensitivity		CD8 α (2.43) Rat mAb (PE-Cy7 $^{\otimes}$ Conjugate) recognizes endogenous levels of total CD8 α protein. This antibody detects an epitope within the extracellular domain.			
Source / Purification		This monoclonal antibody was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation.			
Description		This Cell Signaling Technology antibody is conjugated to PE-Cy7 [®] and tested in-house for direct flow cytometry analysis in mouse cells.			
Background		Cluster of Differentiation 8 (CD8) is a disulphide-linked heterodimer consisting of the unrelated α and β subunits. Each subunit is a glycoprotein composed of a single extracellular Ig-like domain, a polypeptide linker, a transmembrane part and a short cytoplasmic tail. On T cells, CD8 is the coreceptor for the T cell receptor (TCR), and these two distinct structures recognize the Antigen–Major Histocompatibility Complex (MHC). Specifically, the Ig-like domain of CD8 α interacts with the α -domain of the MHC class I molecule. CD8 ensures specificity of the TCR–antigen interaction, prolongs the contact between the T cell and the antigen presenting cell, and the α chain recruits the tyrosine kinase Lck, which is essential for T cell activation (1).			
Background References		1. Zamoyska, R. (1994) <i>Immunity</i> 1, 243-46.			
Species Reactivity		Species reactivity is determined by testing in at least one approved application (e.g., western blot).			
Applications Key		FC-L: Flow Cytometry (Live)			
Cross-Reactivity Key		M: Mouse			
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.			
		Cy and CyDye are registered trademarks of GE Healthcare.			
		All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.			
Limited Uses		Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.			
		Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapoutic purpose, or otherwise in			

purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a

component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.