



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C
#8897

Derlin-1 Antibody

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP	Reactivity: H M R Mk	Sensitivity: Endogenous	MW (kDa): 22	Source/Isotype: Rabbit	UniProt ID: #Q9BUN8	Entrez-Gene Id: 79139
-------------------------------	--------------------------------	-----------------------------------	------------------------	----------------------------------	-------------------------------	---------------------------------

Product Usage Information

Application

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

Derlin-1 Antibody recognizes endogenous levels of total Derlin-1 protein. Based upon sequence alignment, this antibody is not predicted to cross-react with either Derlin-2 or Derlin-3.

Species predicted to react based on 100% sequence homology

Bovine, Dog, Pig, Horse, Guinea Pig

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human Derlin-1 protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

Elimination of misfolded proteins from the endoplasmic reticulum (ER) occurs largely through the ER-associated degradation (ERAD) pathway and is an important physiological adaptation to ER stress. After insertion into the lumen of the ER, glycoproteins that fail to fold properly are destined for degradation. Through a process termed retro-translocation, misfolded proteins are deposited into the cytosol from the ER, where ubiquitination, deglycosylation, and proteasomal proteolysis lead to their degradation. Derlin-1 (Der1-like protein) corresponds to a homologue of yeast Der1p, a protein identified in a genetic screen for components required for the degradation of misfolded ER luminal proteins (1). Like yeast Der1p, mammalian Derlin-1 is an ER protein that is predicted to have four transmembrane segments with both the amino and carboxy termini exposed to the cytoplasmic compartment (2-4). Derlin-1 appears to be a central, evolutionarily conserved membrane component of the retro-translocation machinery associated with the ERAD pathway. Indeed, studies have shown that Derlin-1 expression is transcriptionally upregulated in response to ER stress (5-7) and associates with ER-anchored ubiquitin ligases, such as HRD1 and gp78/AMFR, via binding to p97/VCP and VCP-interacting membrane protein (VIMP) (5,8).

Background References

- Knop, M. et al. (1996) *EMBO J* 15, 753-63.
- Hitt, R. and Wolf, D.H. (2004) *FEMS Yeast Res* 4, 721-9.
- Lilley, B.N. and Ploegh, H.L. (2004) *Nature* 429, 834-40.
- Ye, Y. et al. (2004) *Nature* 429, 841-7.
- Lilley, B.N. and Ploegh, H.L. (2005) *Proc Natl Acad Sci USA* 102, 14296-301.
- Travers, K.J. et al. (2000) *Cell* 101, 249-58.
- Oda, Y. et al. (2006) *J Cell Biol* 172, 383-93.
- Ye, Y. et al. (2005) *Proc Natl Acad Sci USA* 102, 14132-8.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.