Cell Signaling Store at -20C **Filamin B Antibody** H. 877-616-CELL (2355) orders@cellsignal.com Orders: Support: 877-678-TECH (8324) 8949 info@cellsignal.com cellsignal.com Web: 3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H M Mk	Sensitivity: Endogenous	MW (kDa): 280	Source/Isotype: Rabbit	UniProt ID: #075369	Entrez-Gene Id: 2317
Product Usage Information		Application Western Blotting			Dilution 1:1000	
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.				
Specificity/Sensitivity		Filamin B Antibody recognizes endogenous levels of total filamin B protein. This antibody detects a 100 kDa background band of unknown origin. Based on the amino acid sequence, this antibody is not expected to cross-react with filamin A or filamin C.				
Source / Purification		Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human filamin B protein. Antibodies are purified by protein A and peptide affinity chromatography.				
Background		Filamins are a family of dimeric actin binding proteins that function as structural components of cell adhesion sites. They also serve as a scaffold for subcellular targeting of signaling molecules (1). The actin binding domain (α -actinin domain) located at the amino terminus is followed by as many as 24 tandem repeats of about 96 residues and the dimerization domain is located at the carboxy terminus. In addition to actin filaments, filamins associate with other structural and signaling molecules such as β -integrins, Rho/Rac/Cdc42, PKC and the insulin receptor, primarily through the carboxy-terminal dimerization domain (1-3). Filamin A, the most abundant, and filamin B are widely expressed isoforms, while filamin C is predominantly expressed in muscle (1). Filamin A is phosphorylated by PAK1 at Ser2152, which is required for PAK1-mediated actin cytoskeleton reorganization (4).				
Background References		1. Stossel, T.P. et al. (2001) <i>Nat Rev Mol Cell Biol</i> 2, 138-45. 2. Tigges, U. et al. (2003) <i>J Biol Chem</i> 278, 23561-9. 3. He, H.J. et al. (2003) <i>J Biol Chem</i> 278, 27096-104. 4. Vadlamudi, R.K. et al. (2002) <i>Nat Cell Biol</i> 4, 681-90.				
Species Reactivity		Species reactivity is determined by testing in at least one approved application (e.g., western blot).				
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.				
Applications Key		W: Western Blotting				
Cross-Reactivity Key		H: Human M: Mouse Mk: Monkey				
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.				
		All other trademarks a more information.	are the property of t	heir respective owners.	Visit cellsignal.com	/trademarks for
Limited Uses		Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.				
		approved, cleared, or purpose. Customer sh	licensed by the FDA nall not use any Pro-	se Only or a similar labe or other regulatory for duct for any diagnostic o g statement. Products so	eign or domestic er or therapeutic purp	ntity, for any ose, or otherwise in

Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.