

## 8868

## DDX6/RCK (D26C11) Rabbit mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

## For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W	H M R Mk	Endogenous	54	Rabbit IgG	#P26196	1656
Product Usage Information	•	<b>Application</b> Western Blotting			<b>Dilution</b> 1:1000	
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.				
Specificity/Sensitivity		DDX6/RCK (D26C11) Rabbit mAb recognizes endogenous levels of total DDX6/RCK protein.				
Species predicted to react based on 100% sequence homology		Xenopus				
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ala130 of human DDX6/RCK protein.				
Background		DDX6, also known as RCK and p54, was identified as a proto-oncogene product and is a member of the ATP-dependent DEAD box helicase family (1,2). This protein interacts with translation initiation factor eIF4E in the cytoplasmic P-bodies (3) and represses mRNA translation (4). DDX6 is a component of the miRNA induced silencing complex (miRISC) and interacts with Argonaute 1 (Ago1) and Argonaute 2 (Ago2) proteins <i>in vitro</i> and <i>in vivo</i> (5), functioning in miRNA-mediated translational repression (5). Depletion of DDX6 leads to the disruption of cytoplasmic P-bodies indicating that it is required for P-body formation (5).				
Background References		1. Akao, Y. et al. (1995) <i>Cancer Res</i> 55, 3444-9. 2. Weston, A. and Sommerville, J. (2006) <i>Nucleic Acids Res</i> 34, 3082-94. 3. Andrei, M.A. et al. (2005) <i>RNA</i> 11, 717-27. 4. Minshall, N. and Standart, N. (2004) <i>Nucleic Acids Res</i> 32, 1325-34. 5. Chu, C.Y. and Rana, T.M. (2006) <i>PLoS Biol</i> 4, e210.				
Species Reactivity		Species reactivity is determined by testing in at least one approved application (e.g., western blot).				
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.				
Applications Key		W: Western Blotting				
Cross-Reactivity Key		H: Human M: Mouse R: Rat Mk: Monkey				
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.				
		All other trademarks a more information.	are the property of	their respective owners.	Visit cellsignal.com	/trademarks for
Limited Uses		Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.				

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for

Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.