#9147 Store at -20C

MEK2 (13E3) Rabbit mAb	Ce T E			
	Orders:	877-616-CELL (2355) orders@cellsignal.com		
	Support:	877-678-TECH (8324)		
	Web:	info@cellsignal.com		

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H M R Mk	Sensitivity: Endogenous	MW (kDa): 45	Source/Isotype: Rabbit IgG	UniProt ID: #P36507	Entrez-Gene Id: 5605	
Product Usage Information	2	Application Western Blotting			Dilution 1:1000		
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.					
Specificity/Ser	pecificity/Sensitivity MEK2 (13E3) Rabbit mAb detects endogenous levels of total MEK2 protein. This antibody does cross-react with MEK1 and other MAP kinase kinases.		ody does not				
Source / Purifi	cation	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to human MEK2.				prresponding to	
Background		MEK1 and MEK2, also called MAPK or Erk kinases, are dual-specificity protein kinases that function in a mitogen activated protein kinase cascade controlling cell growth and differentiation (1-3). Activation of MEK1 and MEK2 occurs through phosphorylation of two serine residues at positions 217 and 221, located in the activation loop of subdomain VIII, by Raf-like molecules. MEK1/2 is activated by a wide variety of growth factors and cytokines and also by membrane depolarization and calcium influx (1-4). Constitutively active forms of MEK1/2 are sufficient for the transformation of NIH/3T3 cells or the differentiation of PC-12 cells (4). MEK activates p44 and p42 MAP kinase by phosphorylating both threonine and tyrosine residues at sites located within the activation loop of kinase subdomain VIII.					
Background R	eferences	1. Crews, C.M. et al. (1 2. Alessi, D.R. et al. (1 3. Rosen, L.B. et al. (19 4. Cowley, S. et al. (19	994) <i>EMBO J.</i> 13, 16 994) <i>Neuron</i> 12, 12	10-19.			
Species Reacti	vity	Species reactivity is de	etermined by testin	g in at least one approve	ed application (e.g.,	western blot).	
Western Blot B	Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.					
Applications K	ey	W: Western Blotting					
Cross-Reactivi	ty Key	H: Human M: Mouse	R: Rat Mk: Monkey				
Trademarks ar	nd Patents	Cell Signaling Technol	logy is a trademark	of Cell Signaling Techno	ology, Inc.		
		All other trademarks a more information.	are the property of	their respective owners.	Visit cellsignal.com	/trademarks for	
Limited Uses		the following terms a terms and conditions	pply to Products pr that are in additior	a writing signed by a leg ovided by CST, its affiliat o to, or different from, th y authorized representa	es or its distributors lose contained here	s. Any Customer's in, unless	
		approved, cleared, or purpose. Customer sh any manner that conf Customer as the end- diagnostic, prophylac	licensed by the FD/ nall not use any Pro licts with its labelin user and solely for tic or therapeutic p	Jse Only or a similar labe A or other regulatory for duct for any diagnostic o g statement. Products so research and developmo urposes, or any purchas se, requires a separate lig	eign or domestic en or therapeutic purp old or licensed by C ent uses. Any use of e of Product for res	itity, for any ose, or otherwise in ST are provided for Product for ale (alone or as a	

sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.