

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C
#9380

Phospho-RanBP3 (Ser58) Antibody

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP	H Mk	Endogenous	70	Rabbit	#Q9H6Z4	8498

Product Usage Information

Application

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

Phospho-RanBP3 (Ser58) Antibody recognizes endogenous levels of RanBP3b protein only when phosphorylated at Ser58, and RanBP3a protein only when phosphorylated at corresponding residue Ser126.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ser58 of human RanBP3b protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

RanBP3 was originally identified as RanGTP binding protein located in the nucleus and involved in the nuclear exporting process (1). It functions as a cofactor for CRM1 nuclear export by binding to CRM1, stabilizing the RanGTP-CRM1-cargo interaction and promoting complex association with nuclear pore proteins (2,3). In the absence of Ran-bound GTP, RanBP3 prevents binding of CRM1 complex to the nuclear pore complex. In addition to CRM1, RanBP3 also has been shown to bind to RanGEF-RCC1 and increase the guanine nucleotide exchange activity of RCC1 for RanGTP-CRM1-Cargo (1,4). In some cases, as with β-catenin and Smad2/3, RanBP3 binding may mediate the target protein nuclear export in a Ran-dependent, but CRM1-independent manner (5,6). RanBP3 is phosphorylated at Ser58 through the PI3K/Akt or ERK/RSK pathway. This phosphorylation is important for RanBP3 function in nuclear export, likely due to stimulation of RCC1 activity (7,8).

Background References

- Mueller, L. et al. (1998) *FEBS Lett* 427, 330-6.
- Lindsay, M.E. et al. (2001) *J Cell Biol* 153, 1391-402.
- Englmeier, L. et al. (2001) *EMBO Rep* 2, 926-32.
- Nemergut, M.E. et al. (2002) *J Biol Chem* 277, 17385-8.
- Hendriksen, J. et al. (2005) *J Cell Biol* 171, 785-97.
- Dai, F. et al. (2009) *Dev Cell* 16, 345-57.
- Yoon, S.O. et al. (2008) *Mol Cell* 29, 362-75.
- von Knethen, A. et al. (2010) *J Cell Sci* 123, 192-201.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: Human **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless

separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com
For Research Use Only. Not for Use in Diagnostic Procedures.