

Phospho-CDK Substrate Motif [(K/H)pSP] MultiMab[®] Rabbit mAb mix



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures

Applications: W, IP, E-P	Reactivity: All	Sensitivity: Endogenous	Source/Isotype: Rabbit IgG	
Product Usage Information		Application		Dilution
		Western Blotting		1:1000
		Immunoprecipitation		1:50
		Peptide ELISA (DELFIA)		1:1000
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.		
Specificity/Sensitivity		Phospho-CDK Substrate Motif [(K/H)pSP] MultiMab [®] Rabbit mAb mix recognizes phospho-serine in a (K/H)S*P motif. The antibody does not cross-react with phospho-threonine or phospho-tyrosine containing peptides/proteins.		
Source / Purification		MultiMab [®] rabbit monoclonal mix antibodies are prepared by combining individual rabbit monoclonal clones in optimized ratios for the approved applications. Each antibody in the mix is carefully selected based on motif recognition and performance in multiple assays. Each mix is engineered to yield the broadest possible coverage of the modification being studied while ensuring a high degree of specificity for the modification or motif.		
Background		Cyclin-dependent kinases (CDKs) are a family of Ser/Thr kinases that regulate cell-cycle transitions through their association and subsequent phosphorylation of targets in a strictly ordered fashion (1). The substrates for CDKs are proline-directed. The consensus amino acid sequence for CDK substrate is (K/R)(S*)PX(K/R), where X denotes any one of the 20 amino acids (2-4) and S* is the phosphorylation site. Phospho-CDK Substrate Motif [(K/H)pSP] MultiMab™ Rabbit mAb mix recognizes phosphorylated CDK substrates at their consensus motif, providing a powerful tool for CDK target discovery and characterization, as well as HTS drug screening for potential kinase regulators.		
Background References		1. Morgan, D.O. (1997) <i>Annu Rev Cell Dev Biol</i> 13, 261-91. 2. Songyang, Z. et al. (1996) <i>Mol Cell Biol</i> 16, 6486-93. 3. Songyang, Z. (1999) <i>Prog Biophys Mol Biol</i> 71, 359-72. 4. Holmes, J.K. and Solomon, M.J. (1996) <i>J Biol Chem</i> 271, 25240-6.		
Species Reactivi		Species reactivity is deter	mined by testing in at least one a	pproved application (e.g., western blot).
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.		

Applications Key

W: Western Blotting IP: Immunoprecipitation E-P: Peptide ELISA (DELFIA)

Cross-Reactivity Key

All: All Species Expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for their respective owners are the property of their respective owners. Visit cellsignal.com/trademarks for their respective owners are the property of their respective owners. Visit cellsignal.com/trademarks for their respective owners are the property of their respective owners. Visit cells are the property of their respective owners are the property of their respective owners. Visit cells are the property of their respective owners are the property of their respective owners. Visit cells are the property of their respective owners are the property of their respective owners are the property of their respective owners. Visit cells are the property of the

more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.