



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C
#9509

Cleaved Caspase-9 (Asp353) Antibody

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IF-IC	M	Endogenous	37	Rabbit	#Q8C3Q9	12371

Product Usage Information	Application	Dilution
	Western Blotting	1:1000
	Immunofluorescence (Immunocytochemistry)	1:50
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.	
Specificity/Sensitivity	Cleaved Caspase-9 (Asp353) Antibody detects endogenous levels of the 37kDa subunit of mouse caspase-9 only after after cleavage at aspartic acid 353. It does not cross-react with full length caspase-9 or with other caspases at endogenous levels. Non-specific proteins that are induced by apoptosis under certain conditions may be detected.	
Source / Purification	Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Asp353 of mouse caspase-9. Antibodies are purified by protein A and peptide affinity chromatography.	
Background	Caspase-9 (ICE-LAP6, Mch6) is an important member of the cysteine aspartic acid protease (caspase) family (1,2). Upon apoptotic stimulation, cytochrome c released from mitochondria associates with the 47 kDa procaspase-9/Apaf-1. Apaf-1 mediated activation of caspase-9 involves intrinsic proteolytic processing, resulting in cleavage at Asp315 and producing a p35 subunit. Another cleavage occurs at Asp330, producing a p37 subunit that can serve to amplify the apoptotic response (3-6). Cleaved caspase-9 further processes other caspase members, including caspase-3 and caspase-7, to initiate a caspase cascade, which leads to apoptosis (7-10).	
Background References	<ol style="list-style-type: none"> Duan, H. et al. (1996) <i>J. Biol. Chem.</i> 271, 16720-16724. Srinivasula, S. M. et al. (1996) <i>J. Biol. Chem.</i> 271, 27099-27106. Liu, X. et al. (1996) <i>Cell</i> 86, 147-157. Li, P. et al. (1997) <i>Cell</i> 91, 479-489. Zou, H. et al. (1999) <i>J. Biol. Chem.</i> 274, 11549-11556. Srinivasula, S.M. et al. (1998) <i>Mol Cell</i> 1, 949-57. Deveraux, Q. L. et al. (1998) <i>EMBO J.</i> 17, 2215-2223. Slee, E. A. et al. (1999) <i>J. Cell Biol.</i> 144, 281-292. Sun, X.M. et al. (1999) <i>J Biol Chem</i> 274, 5053-60. MacFarlane, M. et al. (1997) <i>J. Cell Biol.</i> 137, 469-479. 	
Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).	
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.	
Applications Key	W: Western Blotting IF-IC: Immunofluorescence (Immunocytochemistry)	
Cross-Reactivity Key	M: Mouse	
Trademarks and Patents	<p>Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.</p> <p>Alexa Fluor is a registered trademark of Life Technologies Corporation.</p> <p>All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.</p>	
Limited Uses	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's	

terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.