



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C
#9607

Phospho-ATM/ATR Substrate (S*Q) (D23H2/D69H5) MultiMab[®] Rabbit mAb mix

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP	Reactivity: All	Sensitivity: Endogenous	Source/Isotype: Rabbit IgG
-------------------------------	---------------------------	-----------------------------------	--------------------------------------

Product Usage Information	Application	Dilution
	Western Blotting	1:1000
	Immunoprecipitation	1:50
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.	
Specificity/Sensitivity	Phospho-ATM/ATR Substrate Motif (S*Q) (D23H2/D69H5) MultiMab [®] Rabbit mAb mix recognizes peptides and proteins containing sequences of phospho-Ser followed by Gln at the +1 position. The antibody does not cross-react with corresponding non-phosphorylated sequences or with other phospho-Ser containing motifs.	
Source / Purification	MultiMab [®] rabbit monoclonal mix antibodies are prepared by combining individual rabbit monoclonal clones in optimized ratios for the approved applications. Each antibody in the mix is carefully selected based on motif recognition and performance in multiple assays. Each mix is engineered to yield the broadest possible coverage of the modification being studied while ensuring a high degree of specificity for the modification or motif.	
Background	Ataxia telangiectasia mutated kinase (ATM) and ataxia telangiectasia and Rad3-related kinase (ATR) are related kinases that regulate cell cycle checkpoints and DNA repair (1). The identified substrates for ATM are p53, p95/NBS1, MDM2, Chk2, BRCA1, CtIP, 4E-BP1, and Chk1 (1,2) The essential requirement for the substrates of ATM/ATR is S*/T*Q. Hydrophobic amino acids at positions -3 and -1, and negatively charged amino acids at position +1 are positive determinants for substrate recognition by these kinases. Positively charged residues surrounding the S*/T*Q are negative determinants for substrate phosphorylation (3). The complex phenotype of AT cells suggests that it likely has additional substrates (3). To better understand the kinase and identify substrates for ATM and the related kinase ATR, CST has developed antibodies that recognize phosphorylated serine or threonine in the S*/T*Q motif.	
Background References	<ol style="list-style-type: none"> 1. Kastan, M.B. and Lim, D.S. (2000) <i>Nature Rev. Mol. Cell Biol.</i> 1, 179-186. 2. Zhao, H. and Piwnica-Worms, H. (2001) <i>Mol. Cell. Biol.</i> 21, 4129-4139. 3. Kim, S. T. et al. (1999) <i>J. Biol. Chem.</i> 274, 37538-37543. 	

Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.
Applications Key	W: Western Blotting IP: Immunoprecipitation
Cross-Reactivity Key	All: All Species Expected
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. MultiMab is a registered trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.
Limited Uses	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.