

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

#9782 Store at -20C

Epithelial-Mesenchymal Transition (EMT) Antibody Sampler Kit

1 Kit (9 x 20 microliters)

For Research Use Only. Not for Use in Diagnostic Procedures.

Product Includes	Product #	Quantity	Mol. Wt	Isotype/Source
Vimentin (D21H3) XP® Rabbit mAb	5741	20 µl	57 kDa	Rabbit IgG
N-Cadherin (D4R1H) XP® Rabbit mAb	13116	20 µl	140 kDa	Rabbit IgG
Claudin-1 (D5H1D) XP® Rabbit mAb	13255	20 µl	20 kDa	Rabbit IgG
β-Catenin (D10A8) XP® Rabbit mAb	8480	20 µl	92 kDa	Rabbit IgG
ZO-1 (D7D12) Rabbit mAb	8193	20 µl	220 kDa	Rabbit IgG
Snail (C15D3) Rabbit mAb	3879	20 µl	29 kDa	Rabbit IgG
Slug (C19G7) Rabbit mAb	9585	20 µl	30 kDa	Rabbit IgG
ZEB1 (D80D3) Rabbit mAb	3396	20 µl	200 kDa	Rabbit IgG
E-Cadherin (24E10) Rabbit mAb	3195	20 µl	135 kDa	Rabbit IgG
Anti-rabbit IgG, HRP-linked Antibody	7074	100 µl		Goat

Please visit cellsignal.com for individual component applications, species cross-reactivity, dilutions, protocols, and additional product information.**Description**

The Epithelial-Mesenchymal Transition (EMT) Antibody Sampler Kit provides an economical means of evaluating EMT. The kit contains enough primary antibody to perform two western blots per primary.

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Background

Epithelial-mesenchymal transition (EMT) is an essential process during development whereby epithelial cells acquire mesenchymal, fibroblast-like properties and display reduced intracellular adhesion and increased motility. This is a critical feature of normal embryonic development, which is also utilized by malignant epithelial tumors to spread beyond their origin (1-3). This tightly regulated process is associated with a number of cellular and molecular events. EMT depends on a reduction in expression of cell adhesion molecules. Cadherins mediate calcium-dependent cell-cell adhesion and play critical roles in normal tissue development (4). E-cadherin is considered an active suppressor of invasion and growth of many epithelial cancers (4-6). Recent studies indicate that cancer cells have up-regulated N-cadherin in addition to loss of E-cadherin. This change in cadherin expression is called the "cadherin switch" and downregulation of E-cadherin is one of the hallmarks of EMT (1). Tight junctions, or zonula occludens, form a continuous barrier to fluids across the epithelium and endothelium. They function in regulation of paracellular permeability and in the maintenance of cell polarity, blocking the movement of transmembrane proteins between the apical and the basolateral cell surfaces. Tight junctions are composed of claudin and occludin proteins, which join the junctions to the cytoskeleton (7,8). Zona occludens proteins ZO-1, 2, and 3 (also known as TJP 1, 2, and 3) are peripheral membrane adaptor proteins that link junctional transmembrane proteins such as occludin and claudin to the actin cytoskeleton (9). ZO-1 and -2 are required for tight junction formation and function (10,11); mutations in ZO-1 and Claudin induce EMT (12). Vimentin is an intermediate filament of mesenchymal origin and is present at early developmental stages. Vimentin's dynamic structural changes and spatial reorganization in response to extracellular stimuli helps to coordinate various signaling pathways (13). β-catenin is a key downstream effector in the Wnt signaling pathway (14). It is implicated in two major biological processes in vertebrates: early embryonic development (15) and tumorigenesis (16). β-catenin also activates Slug. Slug (SNAI2) is a widely expressed transcriptional repressor and member of the Snail family of zinc finger transcription factors (17). Similar to the related Snail protein, Slug binds to the E-cadherin promoter region to repress transcription during development (18). The binding of Slug to integrin promoter sequences represses integrin expression and results in reduced cell adhesion (19). Down regulation of E-cadherin expression occurs during the EMT during embryonic development (20). ZEB family proteins are zinc finger and homeobox domain containing transcription factors. One of the targets suppressed by ZEB proteins is E-cadherin (1).

Background References

1. Aigner, K. et al. (2007) *Oncogene* 26, 6979-88.
2. Peinado, H. et al. (2007) *Nat Rev Cancer* 7, 415-28.

3. Moreno-Bueno, G. et al. (2008) *Oncogene* 27, 6958-69.
 4. Wheelock, M.J. and Johnson, K.R. (2003) *Annu Rev Cell Dev Biol* 19, 207-35.
 5. Christofori, G. (2003) *EMBO J* 22, 2318-23.
 6. Hazan, R.B. et al. (2004) *Ann N Y Acad Sci* 1014, 155-63.
 7. Shin, K. et al. (2006) *Annu Rev Cell Dev Biol* 22, 207-35.
 8. Oliveira, S.S. and Morgado-Díaz, J.A. (2007) *Cell Mol Life Sci* 64, 17-28.
 9. Matter, K. and Balda, M.S. (2007) *J Cell Sci* 120, 1505-11.
 10. Hernandez, S. et al. (2007) *Exp Cell Res* 313, 1533-47.
 11. Umeda, K. et al. (2006) *Cell* 126, 741-54.
 12. Reichert, M. et al. (2000) *J Biol Chem* 275, 9492-500.
 13. Helfand, B.T. et al. (2004) *J Cell Sci* 117, 133-41.
 14. Cadigan, K.M. and Nusse, R. (1997) *Genes Dev* 11, 3286-305.
 15. Wodarz, A. and Nusse, R. (1998) *Annu Rev Cell Dev Biol* 14, 59-88.
 16. Polakis, P. (1999) *Curr Opin Genet Dev* 9, 15-21.
 17. Inukai, T. et al. (1999) *Mol Cell* 4, 343-52.
 18. Bolós, V. et al. (2003) *J Cell Sci* 116, 499-511.
 19. Turner, F.E. et al. (2006) *J Biol Chem* 281, 21321-31.
 20. Barrallo-Gimeno, A. and Nieto, M.A. (2005) *Development* 132, 3151-61.
-

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.