

Revision 1

Store at  
-20C  
#97845**ZMYND8 Antibody**
**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP	H Mk	Endogenous	180	Rabbit	#Q9ULU4	23613

**Product Usage Information****Application**Western Blotting  
Immunoprecipitation**Dilution**1:1000  
1:50**Storage**

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

**Specificity/Sensitivity**

ZMYND8 Antibody recognizes endogenous levels of total ZMYND8 protein.

**Source / Purification**

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues near the amino terminus of human ZMYND8 protein. Antibodies are purified by protein A and peptide affinity chromatography.

**Background**

Zinc finger MYND domain-containing protein 8 (ZMYND8), also referred to as receptor for activated C-kinase 7 (Rack7) and protein kinase C-binding protein 1 (PRKCBP1), is a DNA damage response protein and a transcriptional regulator that is a close homolog of ZMYND11 (BS69) (1). ZMYND8 binds to H3K36me2 and H4K16ac, two histone marks associated with active transcription (2). This protein is targeted to sites of DNA damage within actively transcribed genes, and recruits the H3K4me3-specific histone demethylase KDM5A/JARID1A and nucleosome remodeling and histone deacetylation (NuRD) complex (1-3). Together, these protein complexes mediate transcriptional repression and allow for subsequent double-strand break repair via homologous recombination. ZMYND8 contains a bromodomain and a PWWP domain near its N-terminus, and a MYND domain towards the C-terminus, the latter of which mediates interaction with the NuRD complex (1). ZMYND8 also functions to recruit the H3K4me3-specific histone demethylase KDM5C/JARID1C to enhancer and super-enhancer regions, and functions as a negative regulator of gene expression (4). ZMYND8 and JARID1C are both putative tumor suppressor proteins, and knockdown of either of these proteins leads to derepression of S100 oncogenes (1). ZMYND8 expression is altered in breast and cervical cancer (4,5), and has been found to be translocated with RELA in at least one patient with acute erythroid leukemia (6). Knockdown of ZMYND8 expression in breast cancer cell lines increases anchorage-independent cell growth, cell migration and invasion, and tumor growth in mouse xenograft models (4).

**Background References**

- Gong, F. et al. (2015) *Genes Dev* 29, 197-211.
- Adhikary, S. et al. (2016) *J Biol Chem* 291, 2664-81.
- Gong, F. et al. (2017) *J Cell Biol* 216, 1959-74.
- Shen, H. et al. (2016) *Cell* 165, 331-42.
- Bierkens, M. et al. (2013) *Genes Chromosomes Cancer* 52, 56-68.
- Panagopoulos, I. et al. (2013) *PLoS One* 8, e63663.

**Species Reactivity**

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

**Western Blot Buffer**

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

**Applications Key****W:** Western Blotting **IP:** Immunoprecipitation**Cross-Reactivity Key****H:** Human **Mk:** Monkey**Trademarks and Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.**Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless

separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

**Orders: 877-616-CELL (2355) • [orders@cellsignal.com](mailto:orders@cellsignal.com) • Support: 877-678-TECH (8324) • [info@cellsignal.com](mailto:info@cellsignal.com) • Web: [cellsignal.com](http://cellsignal.com)**  
For Research Use Only. Not for Use in Diagnostic Procedures.