Zap-70 Blocking Peptide



Orders 877-616-CELL (2355)

orders@cellsignal.com

Support 877-678-TECH (8324)

info@cellsignal.com

Web www.cellsignal.com

rev. 05/25/17

For Research Use Only. Not For Use In Diagnostic Procedures.

Description: This peptide can be used to block Zap-70 (99F2) Rabbit mAb #2705.

Background: The Syk family protein tyrosine kinase Zap-70 is expressed in T and NK cells and plays a critical role in mediating T cell activation in response to T cell receptor (TCR) engagement (1). Following TCR engagement, Zap-70 is rapidly phosphorylated on several tyrosine residues through autophosphorylation and transphosphorylation by the Src family tyrosine kinase Lck (2-6). Tyrosine phosphorylation correlates with increased Zap-70 kinase activity and downstream signaling events. Expression of Zap-70 is correlated with disease progression and survival in patients with chronic lymphocytic leukemia (7,8).

Quality Control: The quality of the peptide was evaluated by reversed-phase HPLC and by mass spectrometry. The peptide blocks Zap-70 (99F2) Rabbit mAb #2705 signal in peptide dot blot.

Directions for Use: Use as a blocking reagent to evaluate the specificity of antibody reactivity in peptide dot blot protocols

Entrez Gene ID # UniProt ID

Storage: Supplied in 20 mM potassium phosphate (pH 7.0), 50 mM NaCl, 0.1 mM EDTA, 1 mg/ml BSA and 5% glycerol. 1% DMSO Store at -20°C.

For product specific protocols please see the web page for this product at www.cellsignal.com.

Please visit www.cellsignal.com for a complete listing of recommended complementary products.

Background References:

- (1) Chu, D.H. et al. (1998) Immunol. Rev. 165, 167-180.
- (2) Iwashima, M. et al. (1994) Science 263, 1136-1139.
- (3) Neumeister, E.N. et al. (1995) Mol. Cell Biol. 15, 3171-3178.
- (4) Chan, A.C. et al. (1995) EMBO J. 14, 2499-2508.
- (5) Williams, B.L. et al. (1999) EMBO J. 18, 1832-1844.
- (6) Di Bartolo, V. et al. (1999) J. Biol. Chem. 274, 6285-6294.
- (7) Wiestner, A. et al. (2003) Blood 101, 4944-4951.
- (8) Crespo, M. et al. (2003) N. Engl. J. Med. 348, 1764-1775.