## **IFIT1 Antibody**



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

## For Research Use Only. Not for Use in Diagnostic Procedures.

<b>Applications:</b> W, IP	Reactivity: H	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 56	Source/Isotype: Rabbit	<b>UniProt ID:</b> #P09914	Entrez-Gene Id: 3434	
Product Usage		Application			Dilution		
Information		Western Blotting			1:1000		
		Immunoprecipitation			1:50		
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 $\mu$ g/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.					
Specificity/Sensitivity		IFIT1 Antibody recognizes endogenous levels of total IFIT1 protein.					
Source / Purification		Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ala194 of human IFIT1 protein. Antibodies are purified by protein A and peptide affinity chromatography.					
Background		IFIT1 (interferon-induced protein with tetratricopeptide repeats 1) belongs to the IFIT family of proteins, which consists of four members in humans (IFIT1, IFIT2, IFIT3, and IFIT5) and three members in mice (IFIT1, IFIT2, and IFIT3) (1). IFIT1 expression is induced by type I interferons resulting from viral infection (2). IFIT1 is an antiviral protein that directly binds viral RNA that has a 5' triphosphate group (PPP-RNA) (2). In humans, the viral PPP-RNA bound IFIT1 forms a complex with IFIT2 and IFIT3 to sequester the viral PPP-RNA and prevent replication (2). IFIT1 has also been shown to inhibit translation by binding to the eukaryotic initiation factor 3 (eIF3) (3).					
Background References		<ol> <li>Ablasser, A. and Hornung, V. (2011) <i>Nat Immunol</i> 12, 588-90.</li> <li>Pichlmair, A. et al. (2011) <i>Nat Immunol</i> 12, 624-30.</li> <li>Guo, J. et al. (2000) <i>EMBO J</i> 19, 6891-9.</li> </ol>					
Species Reacti	vity	Species reactivity is de	termined by testin	g in at least one approve	ed application (e.g.,	western blot).	
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.					

Applications Key

W: Western Blotting IP: Immunoprecipitation

**Cross-Reactivity Key** 

H: Human

**Trademarks and Patents** 

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

**Limited Uses** 

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not

sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.