

#12307 Store at -20°C

SignalSilence® DNMT3L siRNA I (Mouse Specific)

✓ 10µM in 300 µl (3 nmol)



Orders ■ 877-616-CELL (2355) orders@cellsignal.com
Support ■ 877-678-TECH (8324) info@cellsignal.com
Web ■ www.cellsignal.com

rev. 05/16/16


For Research Use Only. Not For Use In Diagnostic Procedures.

Species Cross-Reactivity: M

Description: SignalSilence® DNMT3L siRNA I (Mouse Specific) from Cell Signaling Technology (CST) allows the researcher to specifically inhibit DNMT3L expression using RNA interference, a method whereby gene expression can be selectively silenced through the delivery of double stranded RNA molecules into the cell. All SignalSilence® siRNA products from CST are rigorously tested in-house and have been shown to reduce target protein expression by western analysis.

Background: Methylation of DNA at cytosine residues in mammalian cells is a heritable, epigenetic modification that is critical for proper regulation of gene expression, genomic imprinting, and development (1,2). Three families of mammalian DNA methyltransferases have been identified: DNMT1, DNMT2, and DNMT3 (1,2). DNMT1 is constitutively expressed in proliferating cells and functions as a maintenance methyltransferase, transferring proper methylation patterns to newly synthesized DNA during replication. DNMT3A and DNMT3B are strongly expressed in embryonic stem cells with reduced expression in adult somatic tissues. DNMT3A and DNMT3B function as *de novo* methyltransferases that methylate previously unmethylated regions of DNA. DNMT2 is expressed at low levels in adult somatic tissues and its inactivation affects neither *de novo* nor maintenance DNA methylation.

DNMT3L is a catalytically inactive regulatory factor for the DNMT3A and DNMT3B *de novo* methyltransferases that is expressed at low levels in embryonic stem cells, testis, ovaries, and thymus (1,2). These *de novo* methyltransferases consist of a heterotetrameric complex containing two molecules of DNMT3L, and either two molecules DNMT3A or DNMT3B (3). DNMT3L contains an amino-terminal ATRX-DNMT3-DNMT3L (ADD) domain and a carboxy-terminal methyltransferase-like domain (4-7). The methyltransferase-like domain binds to DNMT3A and DNMT3B to stimulate catalytic activity by increasing the binding of S-adenosylmethionine and DNA (4,5). The ADD domain recruits the methyltransferase complex to transcriptionally inactive regions of the genome by binding to unmethylated histone H3 Lys4 (6,7).



Western blot analysis of extracts from F9 cells, transfected with 100 nM SignalSilence® Control siRNA (Unconjugated) #6568 (-), SignalSilence® DNMT3L siRNA I (Mouse Specific) (+), or SignalSilence® DNMT3L siRNA II (Mouse Specific) #12308 (+), using DNMT3L Antibody (Mouse Specific) #12309 (upper) or β-Actin (D6A8) Rabbit mAb #8457 (lower). The DNMT3L Antibody (Mouse Specific) confirms silencing of DNMT3L expression, while the β-Actin (D6A8) Rabbit mAb is used as a loading control.

Directions for Use: CST recommends transfection with 100 nM SignalSilence® DNMT3L siRNA I (Mouse Specific) 48 to 72 hours prior to cell lysis. For transfection procedure, follow protocol provided by the transfection reagent manufacturer. Please feel free to contact CST with any questions on use.

Each vial contains the equivalent of 100 transfections, which corresponds to a final siRNA concentration of 100 nM per transfection in a 24-well plate with a total volume of 300 µl per well.

Quality Control: Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid phase extraction. The annealed RNA duplex is further analyzed by mass spectrometry to verify the exact composition of the duplex. Each lot is compared to the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency.


Entrez-Gene ID #29947
Swiss-Prot Acc. #Q9UJW3

Storage: DNMT3L siRNA I (Mouse Specific) is supplied in RNase-free water. Aliquot and store at -20°C.

Please visit www.cellsignal.com for a complete listing of recommended companion products.

Background References:

- (1) Hermann, A. et al. (2004) *Cell. Mol. Life Sci.* 61, 2571–2587.
- (2) Turek-Plewa, J. and Jagodziński, P.P. (2005) *Cell. Mol. Biol. Lett.* 10, 631–647.
- (3) Jia, D. et al. (2007) *Nature* 449, 248–251.
- (4) Holz-Schietinger, C. and Reich, N.O. (2010) *J. Biol. Chem.* 285, 29091–29100.
- (5) Suetake, I. et al. (2004) *J. Biol. Chem.* 279, 27816–27823.
- (6) Ooi, S.K. et al. (2007) *Nature* 448, 714–717.
- (7) Otani, J. et al. (2009) *EMBO Rep.* 10, 1235–1241.



Western blot analysis of extracts from F9 cells, transfected with 100 nM SignalSilence® Control siRNA (Unconjugated) #6568 (-), SignalSilence® DNMT3L siRNA I (Mouse Specific) (+), or SignalSilence® DNMT3L siRNA II (Mouse Specific) #12308 (+), using DNMT3L (E1Y7Q) Rabbit mAb (Mouse Specific) #13451 (upper) or GAPDH (D16H11) XP® Rabbit mAb #5174 (lower). The DNMT3L (E1Y7Q) Rabbit mAb (Mouse Specific) confirms silencing of DNMT3L expression, while the GAPDH (D16H11) XP® Rabbit mAb is used as a loading control.

© 2012 Cell Signaling Technology, Inc. XP®, SignalSilence®, CST™, and Cell Signaling Technology® are trademarks of Cell Signaling Technology, Inc.

Applications Key: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide
Species Cross-Reactivity Key: H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine
Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—Horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.