

## **NKX2.2 Antibody**



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

## For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	<b>Sensitivity:</b> Transfected Only	<b>MW (kDa):</b> 40	<b>Source/Isotype:</b> Rabbit	<b>UniProt ID:</b> #O95096	Entrez-Gene Id: 4821
Product Usage Information		<b>Application</b> Western Blotting			<b>Dilution</b> 1:1000	
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.				
Specificity/Sensitivity		NKX2.2 Antibody recognizes transfected levels of total NKX2.2 protein.				
Species predicted to react based on 100% sequence homology		Monkey				
Source / Purification		Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ala92 of human NKX2.2 protein. Antibodies are purified by protein A and peptide affinity chromatography.				
Background		The NKX family of homeobox genes are known to act as intermediaries in the neural response to Sonic hedgehog signaling during central nervous system (CNS) development (1). NKX2.2 is a member of this family of transcription factors and is necessary for neuroendocrine differentiation in the CNS and pancreas (2,3). NKX2.2 mutant mice die shortly after birth due to incomplete differentiation of insulin-producing pancreatic $\beta$ cells and defects in ventral neural patterning (2,3). According to the research literature, expression of NKX2.2 has also been found in neuroendocrine tumors of the gut, making it a potential marker for the study of gastrointestinal neuroendocrine tumors (4).				
Background References		1. Sander, M. et al. (2000) <i>Genes Dev</i> 14, 2134-9. 2. Sussel, L. et al. (1998) <i>Development</i> 125, 2213-21. 3. Briscoe, J. et al. (1999) <i>Nature</i> 398, 622-7. 4. Wang, Y.C. et al. (2009) <i>Endocr Relat Cancer</i> 16, 267-79.				
Species Reactivity		Species reactivity is determined by testing in at least one approved application (e.g., western blot).				
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.				
Applications Key		W: Western Blotting				
Cross-Reactivity Key		H: Human				
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.				
		All other trademarks a more information.	re the property of	their respective owners.	Visit cellsignal.com	/trademarks for
Limited Uses		Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.				

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for

Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.