

Store at
-20°C
#12315**TRIM25 Antibody**

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W	H	Endogenous	72	Rabbit	#Q14258	7706

Product Usage Information**Application**

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

TRIM25 Antibody recognizes endogenous levels of total TRIM25 protein.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Gln146 of human TRIM25 protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

TRIM25, also termed Estrogen-responsive Finger Protein (EFP), is a member of the tripartite motif-containing (TRIM) family of proteins, characterized by the presence of a RING domain, one or two B-box motifs, and a coiled-coil region (1). TRIM25 was first identified in a search for estrogen-responsive genes (2), and studies have subsequently shown TRIM25 to be overexpressed in many breast cancer tumors (3). A potentially oncogenic role for TRIM25 was suggested by studies showing that suppression of TRIM25 expression inhibited growth of MCF7 cells *in vitro* and in mouse xenograft models (4). Functional studies largely suggest that TRIM25 functions as a ubiquitin E3 or ISG15 E3 ligase. For example, TRIM25 was shown to induce K63-linked ubiquitination of Rig-I, resulting in Rig-I-mediated activation of downstream signaling cascades that drive the host antiviral innate immune response (5). Notably, it was reported that the influenza A virus non-structural protein 1 inhibits TRIM25-mediated ubiquitination of Rig-I, which may have evolved as a mechanism to evade the host innate immune response (6).

Background References

1. Meroni, G. and Diez-Roux, G. (2005) *Bioessays* 27, 1147-57.
2. Inoue, S. et al. (1993) *Proc Natl Acad Sci U S A* 90, 11117-21.
3. Suzuki, T. et al. (2005) *Clin Cancer Res* 11, 6148-54.
4. Ueyama, K. et al. (2010) *Cancer Gene Ther* 17, 624-32.
5. Gack, M.U. et al. (2007) *Nature* 446, 916-920.
6. Gack, M.U. et al. (2009) *Cell Host Microbe* 5, 439-49.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting

Cross-Reactivity Key

H: Human

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.