

Store at
-20C
#12362**DLAT (4A4-B6-C10) Mouse mAb**
Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP, IF-IC	H Mk	Endogenous	69	Mouse IgG1	#P10515	1737

Product Usage Information**Application**

Western Blotting
Immunoprecipitation
Immunofluorescence (Immunocytochemistry)

Dilution

1:1000
1:50
1:50 - 1:100

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

DLAT (4A4-B6-C10) Mouse mAb recognizes endogenous levels of total DLAT protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a recombinant human DLAT protein.

Background

Dihydrolipoamide acetyltransferase (DLAT) transfers an acetyl group from pyruvate to CoA to synthesize acetyl-CoA (1-4). This protein, also known as the E2 subunit of the pyruvate dehydrogenase complex (PDC-E2), has been implicated in the literature as the primary autoantigen in primary biliary cirrhosis (2-5). Antimitochondrial antibodies (AMAs) are likely formed when DLAT is exposed to the immune system in apoptotic cells of the bile duct (3,5). Research studies have shown that in some cases, cosmetics, NSAIDs, chewing gum, acetaminophen, and other compounds could trigger exposure of DLAT in sensitive individuals (3). The presence of AMAs is often detectable before disease diagnosis (4,5). Research studies have also shown that activation of the Toll-like receptor-3 (TLR-3) pathway is involved in the progression from a subclinical to clinical state (4).

Background References

1. Thekkumkara, T.J. et al. (1988) *FEBS Lett* 240, 45-8.
2. Coppel, R.L. et al. (1988) *Proc Natl Acad Sci U S A* 85, 7317-21.
3. Naiyanetr, P. et al. (2011) *J Autoimmun* 37, 209-16.
4. Ambrosini, Y.M. et al. (2011) *Clin Exp Immunol* 166, 110-20.
5. Lleo, A. et al. (2009) *Hepatology* 49, 871-9.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation **IF-IC:** Immunofluorescence (Immunocytochemistry)

Cross-Reactivity Key

H: Human **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not

sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com
For Research Use Only. Not for Use in Diagnostic Procedures.