

Store at RT
#12606

16% Formaldehyde, Methanol-Free

50 ml (5 x 10 ml)



Cell Signaling
TECHNOLOGY®

Support: +1-978-867-2388 (U.S.)
www.cellsignal.com/support

Orders: 877-616-2355 (U.S.)
orders@cellsignal.com


rev. 05/19/16

For Research Use Only. Not For Use In Diagnostic Procedures.

Description: Premium 16% (w/v) Formaldehyde from Cell Signaling Technology is used as a fixative agent for fluorescent immunocytochemical, flow cytometry, and chromatin immunoprecipitation (ChIP) assays. It is methanol-free, prepared from high quality paraformaldehyde, and packaged under an inert atmosphere of nitrogen.

Each 10 ml solution is supplied in an amber glass vial with two access points, offering distinct advantages over pre-scored ampules. The screw cap allows for easy access to large volumes if necessary. To extend the product's shelf life, small volumes should be extracted by piercing the silicone top with a needle and syringe.

Molecular Formula: CH₂O



Molecular Weight: 30.03 g/mol

Storage: Store at room temperature. Each vial is stable for twelve months when left unopened or when material is extracted with a needle and syringe. Use within one month once opened.

Directions for Use: Final working concentrations can vary depending on the application, but is always prepared fresh by diluting in 1X PBS just prior to use. Leftover diluted material should be discarded.

Recommended Formaldehyde Fixative Concentrations:

Immunofluorescence (IF-IC and IF-F): 4%
Flow Cytometry: 2-4%
Chromatin IP: 1%

Thank you for your recent purchase. If you would like to provide a review visit www.cellsignal.com/comments.

www.cellsignal.com

© 2014 Cell Signaling Technology, Inc.

Cell Signaling Technology® is a trademark of Cell Signaling Technology, Inc.

Applications: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide **Species Cross-Reactivity:** H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—Horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.