Cleaved Caspase-3 (Asp175) (D3E9) Rabbit mAb (PE Conjugate)



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Applications: FC-FP	Reactivity: H M	Sensitivity: Endogenous	Source/Isotype: Rabbit IgG	UniProt ID: #P42574	Entrez-Gene Id: 836
Product Usage Information		Application Flow Cytometry (Fixed/P	ermeabilized)		Dilution 1:50
Storage		Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2 mg/ml BSA. Store at 4°C. Do not aliquot the antibodies. Protect from light. Do not freeze.			
Specificity/Sensitivity		Cleaved Caspase-3 (Asp175) (D3E9) Rabbit mAb (PE Conjugate) recognizes endogenous levels of caspase-3 protein only when cleaved at Asp175. This antibody may also detect non-specific caspase substrates. Non-specific labeling may be observed by immunofluorescence in specific sub-types of healthy cells in fixed-frozen tissues (e.g. pancreatic alpha-cells). Nuclear background may be observed in rat and monkey samples.			
Species predicted to react based on 100% sequence homology		Rat, Bovine, Dog, Pig			
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Asp175 of human caspase-3 protein.			
Description		This Cell Signaling Technology antibody is conjugated to phycoerythrin (PE) and tested in-house for direct flow cytometry analysis in human cells. The antibody is expected to exhibit the same species cross-reactivity as the unconjugated Cleaved Caspase-3 (Asp175) (D3E9) Rabbit mAb #9579.			
Background		Caspase-3 (CPP-32, Apopain, Yama, SCA-1) is a critical executioner of apoptosis, as it is either partially or totally responsible for the proteolytic cleavage of many key proteins, such as the nuclear enzyme poly (ADP-ribose) polymerase (PARP) (1). Activation of caspase-3 requires proteolytic processing of its inactive zymogen into activated p17 and p12 fragments. Cleavage of caspase-3 requires the aspartic acid residue at the P1 position (2).			
Background References		1. Fernandes-Alnemri, T. et al. (1994) <i>J Biol Chem</i> 269, 30761-4. 2. Nicholson, D.W. et al. (1995) <i>Nature</i> 376, 37-43.			
Species Reactiv	ity	Species reactivity is dete	rmined by testing in at le	ast one approved ap	plication (e.g., western blot).
Applications Key		FC-FP: Flow Cytometry (Fixed/Permeabilized)			
Cross-Reactivity Key		H: Human M: Mouse			
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.			
		All other trademarks are more information.	the property of their res	pective owners. Visit	cellsignal.com/trademarks for
l imited lises		Event as otherwise eve	ressly agreed in a writing	signed by a legally a	uthorized representative of CST

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for

diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.