

# BAF Complex Antibody Sampler Kit

✓ 1 Kit  
(6 x 20 µl)



**Orders** ■ 877-616-CELL (2355)  
orders@cellsignal.com

**Support** ■ 877-678-TECH (8324)  
info@cellsignal.com

**Web** ■ www.cellsignal.com

rev. 04/19/17

**For Research Use Only. Not For Use In Diagnostic Procedures.**

Products Included	Product #	Quantity	Mol. Wt.	Isotype
ARID1A/BAF250A (D2A8U) Rabbit mAb	12354	20 µl	250 kDa	Rabbit IgG
Brg1 (A52) Antibody	3508	20 µl	220 kDa	Rabbit IgG
BRM (D9E8B) XP® Rabbit mAb	11966	20 µl	200 kDa	Rabbit IgG
SMARCC1/BAF155 (D7F8S) Rabbit mAb	11956	20 µl	155 kDa	Rabbit IgG
SMARCC2/BAF170 (D8O9V) Rabbit mAb	12760	20 µl	162, 170 kDa	Rabbit IgG
SMARCB1/BAF47 (D8M1X) Rabbit mAb	91735	20 µl	44 kDa	Rabbit IgG
Anti-rabbit IgG, HRP-linked Antibody	7074	100 µl		Goat

See [www.cellsignal.com](http://www.cellsignal.com) for individual companion applications, species cross-reactivity, dilutions and additional application protocols.

**Description:** The BAF Complex Antibody Sampler Kit provides an economical means of detecting total protein from the SWI/SNF family members including ARID1A/BAF250A, Brg1, BRM, SMARCC1/BAF155, SMARCC2/BAF170 and SMARCB1/BAF47. The kit contains enough primary antibody to perform two western blots per primary antibody.

**Background:** ATP-dependent chromatin remodeling complexes play an essential role in the regulation of various nuclear processes, such as gene expression, DNA replication, and repair (1,2). The SWI/SNF chromatin remodeling complex consists of more than 10 subunits with a single molecule of the ATPase catalytic subunit BRM or BRG1, but not both. The activities of these two subunits drive the disruption of histone-DNA contacts that lead to changes in accessibility of crucial regulatory elements within chromatin (2-5). The BRM/BRG1 containing SWI/SNF complexes are recruited to target promoters by transcription factors, such as nuclear receptors, p53, RB, and BRCA1 to regulate gene activation, cell growth, the cell cycle, and differentiation processes (1,6-9). BRM and BRG1 are also considered to be tumor suppressors and their expression levels are severely reduced in several cancer cell lines (10-13).

SMARCC1/BAF155, SMARCC2/BAF170, and SMARCB1/BAF47 are members of the core subunits of the SWI/SNF complex, which is necessary for efficient nucleosome remodeling by BRG1 in vitro (14). ARID1A/BAF250A is one of the accessory subunits of the SWI/SNF complex (15). SMARCC1, SMARCB1, and ARID1A are an essential part of the mouse embryonic stem cell specific SWI/SNF complex (esBAF). SMARCC1 is necessary for early embryogenesis, especially proper brain and visceral endoderm development (16-18). SMARCB1 is necessary for early embryogenesis and hepatocyte differentiation (19,20).

**Specificity/Sensitivity:** Each antibody in this kit recognizes endogenous levels of total protein for the specified target and does not cross-react with other family members. ARID1A/BAF250A (D2A8U) Rabbit mAb also cross-reacts with proteins of unknown origin at 65 kDa.

**Source/Purification:** Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Gly1293 of human ARID1A/BAF250A protein, Gly264 of human BRM protein, Gly975 of human SMARCC1/BAF155 protein, Ile818 of human SMARCC2/BAF170 protein, or Leu120 of human SMARCB1/BAF47 protein. Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ala52 of human Brg1 protein. Polyclonal antibodies are purified by Protein A and peptide affinity chromatography.

**Storage:** Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibodies.

**Recommended Antibody Dilutions:**  
Western blotting 1:1000

Please visit [www.cellsignal.com](http://www.cellsignal.com) for validation data and a complete listing of recommended companion products.

## Background References:

- (1) Ho, L. and Crabtree, G.R. (2010) *Nature* 463, 474-84.
- (2) Becker, P.B. and Hörz, W. (2002) *Annu Rev Biochem* 71, 247-73.
- (3) Eberharter, A. and Becker, P.B. (2004) *J Cell Sci* 117, 3707-11.
- (4) Bowman, G.D. (2010) *Curr Opin Struct Biol* 20, 73-81.
- (5) Gangaraju, V.K. and Bartholomew, B. (2007) *Mutat Res* 618, 3-17.
- (6) Lessard, J.A. and Crabtree, G.R. (2010) *Annu Rev Cell Dev Biol* 26, 503-32.
- (7) Moretini, S. et al. (2008) *Front Biosci* 13, 5522-32.
- (8) Wolf, I.M. et al. (2008) *J Cell Biochem* 104, 1580-6.
- (9) Simone, C. (2006) *J Cell Physiol* 207, 309-14.
- (10) Yamamichi, N. et al. (2005) *Oncogene* 24, 5471-81.
- (11) Reisman, D.N. et al. (2002) *Oncogene* 21, 1196-207.
- (12) Shen, H. et al. (2008) *Cancer Res* 68, 10154-62.
- (13) Weissman, B. and Knudsen, K.E. (2009) *Cancer Res* 69, 8223-30.
- (14) Phelan, M.L. et al. (1999) *Mol Cell* 3, 247-53.
- (15) Gao, X. et al. (2008) *Proc Natl Acad Sci U S A* 105, 6656-61.
- (16) Han, D. et al. (2008) *Dev Biol* 315, 136-46.
- (17) Kim, J.K. et al. (2001) *Mol Cell Biol* 21, 7787-95.
- (18) Schaniel, C. et al. (2009) *Stem Cells* 27, 2979-91.
- (19) Klochendler-Yeivin, A. et al. (2000) *EMBO Rep* 1, 500-6.
- (20) Gresh, L. et al. (2005) *EMBO J* 24, 3313-24.
- (21) Ho, L. et al. (2009) *Proc Natl Acad Sci USA* 106, 5181-6.

**Applications Key:** W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide

**Species Cross-Reactivity Key:** H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine

Dg—dog Pg—pig Sc—S. cerevisiae All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.

## Western Immunoblotting Protocol

For western blots, incubate membrane with diluted primary antibody in either 5% w/v BSA or nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

**NOTE:** Please refer to primary antibody datasheet or product webpage for recommended primary antibody dilution buffer and recommended antibody dilution.

### A. Solutions and Reagents

**NOTE:** Prepare solutions with reverse osmosis deionized (RODI) or equivalent grade water.

- 20X Phosphate Buffered Saline (PBS):** (#9808) To prepare 1 L 1X PBS: add 50 ml 20X PBS to 950 ml dH<sub>2</sub>O, mix.
- 10X Tris Buffered Saline (TBS):** (#12498) To prepare 1 L 1X TBS: add 100 ml 10X to 900 ml dH<sub>2</sub>O, mix.
- 1X SDS Sample Buffer:** Blue Loading Pack (#7722) or Red Loading Pack (#7723)  
Prepare fresh 3X reducing loading buffer by adding 1/10 volume 30X DTT to 1 volume of 3X SDS loading buffer. Dilute to 1X with dH<sub>2</sub>O.
- 10X Tris-Glycine SDS Running Buffer:** (#4050) To prepare 1 L 1X running buffer: add 100 ml 10X running buffer to 900 ml dH<sub>2</sub>O, mix.
- 10X Tris-Glycine Transfer Buffer:** (#12539) To prepare 1 L 1X transfer buffer: add 100 ml 10X transfer buffer to 200 ml methanol + 700 ml dH<sub>2</sub>O, mix.
- 10X Tris Buffered Saline with Tween® 20 (TBST):** (#9997) To prepare 1 L 1X TBST: add 100 ml 10X TBST to 900 ml dH<sub>2</sub>O, mix.
- Nonfat Dry Milk:** (#9999)
- Blocking Buffer:** 1X TBST with 5% w/v nonfat dry milk; for 150 ml, add 7.5 g nonfat dry milk to 150 ml 1X TBST and mix well.
- Wash Buffer:** (#9997) 1X TBST
- Bovine Serum Albumin (BSA):** (#9998)
- Primary Antibody Dilution Buffer:** 1X TBST with 5% BSA or 5% nonfat dry milk as indicated on primary antibody datasheet; for 20 ml, add 1.0 g BSA or nonfat dry milk to 20 ml 1X TBST and mix well.
- Biotinylated Protein Ladder Detection Pack:** (#7727)
- Prestained Protein Marker, Broad Range (Premixed Format):** (#7720)
- Blotting Membrane and Paper:** (#12369) This protocol has been optimized for nitrocellulose membranes. Pore size 0.2 µm is generally recommended.
- Secondary Antibody Conjugated to HRP:** anti-rabbit (#7074); anti-mouse (#7076)
- Detection Reagent:** LumiGLO® chemiluminescent reagent and peroxide (#7003) or SignalFire™ ECL Reagent (#6883)

### B. Protein Blotting

**A general protocol for sample preparation.**

- Treat cells by adding fresh media containing regulator for desired time.
- Aspirate media from cultures; wash cells with 1X PBS; aspirate.
- Lyse cells by adding 1X SDS sample buffer (100 µl per well of 6-well plate or 500 µl for a 10 cm diameter plate). Immediately scrape the cells off the plate and transfer the extract to a microcentrifuge tube. Keep on ice.
- Sonicate for 10–15 sec to complete cell lysis and shear DNA (to reduce sample viscosity).
- Heat a 20 µl sample to 95–100°C for 5 min; cool on ice.
- Microcentrifuge for 5 min.
- Load 20 µl onto SDS-PAGE gel (10 cm x 10 cm). **NOTE:** Loading of prestained molecular weight markers (#7720, 10 µl/lane) to verify electrotransfer and biotinylated protein ladder (#7727, 10 µl/lane) to determine molecular weights are recommended.
- Electrotransfer to nitrocellulose membrane (#12369).

### C. Membrane Blocking and Antibody Incubations

**NOTE:** Volumes are for 10 cm x 10 cm (100 cm<sup>2</sup>) of membrane; for different sized membranes, adjust volumes accordingly.

#### I. Membrane Blocking

- (Optional) After transfer, wash nitrocellulose membrane with 25 ml TBS for 5 min at room temperature.
- Incubate membrane in 25 ml of blocking buffer for 1 hr at room temperature.
- Wash three times for 5 min each with 15 ml of TBST.

#### II. Primary Antibody Incubation

- Incubate membrane and primary antibody (at the appropriate dilution and diluent as recommended in the product datasheet) in 10 ml primary antibody dilution buffer with gentle agitation overnight at 4°C.
- Wash three times for 5 min each with 15 ml of TBST.
- Incubate membrane with the species appropriate HRP-conjugated secondary antibody (#7074 or #7076 at 1:2000) and anti-biotin, HRP-linked Antibody (#7075 at 1:1000–1:3000) to detect biotinylated protein markers in 10 ml of blocking buffer with gentle agitation for 1 hr at room temperature.
- Wash three times for 5 min each with 15 ml of TBST.
- Proceed with detection (Section D).

### D. Detection of Proteins

- Incubate membrane with 10 ml LumiGLO® (0.5 ml 20X LumiGLO® #7003, 0.5 ml 20X peroxide, and 9.0 ml purified water) or 10 ml SignalFire™ #6883 (5 ml Reagent A, 5 ml Reagent B) with gentle agitation for 1 min at room temperature.
- Drain membrane of excess developing solution (do not let dry), wrap in plastic wrap and expose to x-ray film. An initial 10 sec exposure should indicate the proper exposure time.  
**NOTE:** Due to the kinetics of the detection reaction, signal is most intense immediately following incubation and declines over the following 2 hr.

LumiGLO® is a registered trademark of Kirkegaard & Perry Laboratories. Tween® is a registered trademark of ICI Americas, INC. SignalFire™ is a trademark of Cell Signaling Technology, INC.

Orders ■ 877-616-CELL (2355) orders@cellsignal.com Support ■ 877-678-TECH (8324) info@cellsignal.com Web ■ www.cellsignal.com