

**Phospho-Topoisomerase II $\alpha$  (Ser1469)  
(D4F5) Rabbit mAb****Orders:** 877-616-CELL (2355)  
orders@cellsignal.com**Support:** 877-678-TECH (8324)**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

<b>Applications:</b> W, IF-IC	<b>Reactivity:</b> H	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 190	<b>Source/Isotype:</b> Rabbit IgG	<b>UniProt ID:</b> #P11388	<b>Entrez-Gene Id:</b> 7153
----------------------------------	-------------------------	-----------------------------------	-------------------------	--------------------------------------	-------------------------------	--------------------------------

**Product Usage Information****Application**Western Blotting  
Immunofluorescence (Immunocytochemistry)**Dilution**1:1000  
1:400**Storage**Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100  $\mu$ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at  $-20^{\circ}\text{C}$ . Do not aliquot the antibody.**Specificity/Sensitivity**Phospho-Topoisomerase II $\alpha$  (Ser1469) (D4F5) Rabbit mAb recognizes endogenous levels of topoisomerase II $\alpha$  protein only when phosphorylated at Ser1469.**Source / Purification**Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ser1469 of human topoisomerase II $\alpha$  protein.**Background**

DNA topoisomerases I and II are nuclear enzymes; type II consists of two highly homologous isoforms: topoisomerase II $\alpha$  and II $\beta$ . These enzymes regulate the topology of DNA, maintain genomic integrity, and are essential for processes such as DNA replication, recombination, transcription, and chromosome segregation by allowing DNA strands to pass through each other (1). Topoisomerase I nicks and rejoins one strand of the duplex DNA, while topoisomerase II transiently breaks and closes double-stranded DNA (2). Topoisomerases are very susceptible to various stresses. Acidic pH or oxidative stress can convert topoisomerases to DNA-breaking nucleases, causing genomic instability and cell death. DNA-damaging topoisomerase targeting drugs (e.g., etoposide) also convert topoisomerases to nucleases, with the enzyme usually trapped as an intermediate that is covalently bound to the 5' end of the cleaved DNA strand(s). Research studies have shown that this intermediate leads to genomic instability and cell death. Thus, agents that target topoisomerases are highly sought after cancer chemotherapeutic drugs (3).  $\text{Ca}^{2+}$ -regulated phosphorylation of topoisomerase II $\alpha$  at Ser1106 modulates the activity of this enzyme and its sensitivity to targeting drugs (4). Casein kinase 2 (CK2) phosphorylates DNA topoisomerase II $\alpha$  on Ser1469 to generate a site recognized by the mitosis-specific antibody MPM-2 (5).

**Background References**

1. Wang, J.C. (2002) *Nat. Rev. Mol. Cell. Biol.* 3, 430-40.
2. Pulleyblank, .E. (1997) *Science* 277, 648-9.
3. Li, T.K. and Liu, L.F. (2001) *Annu. Rev. Pharmacol. Toxicol.* 41, 53-77.
4. Chikamori, K. et al. (2003) *J. Biol. Chem.* 278, 12696-702.
5. Escargueil, A.E. et al. (2000) *J Biol Chem* 275, 34710-8.

**Species Reactivity**

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

**Western Blot Buffer****IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at  $4^{\circ}\text{C}$  with gentle shaking, overnight.**Applications Key****W:** Western Blotting **IF-IC:** Immunofluorescence (Immunocytochemistry)**Cross-Reactivity Key****H:** Human**Trademarks and Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.**Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's

terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.