

VAMP1 Antibody



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP	Reactivity: H M R	Sensitivity: Endogenous	MW (kDa): 16	Source/Isotype: Rabbit	UniProt ID: #P23763	Entrez-Gene Id: 6843
Product Usage Information	2	Application Western Blotting Immunoprecipitation			Dilution 1:1000 1:50	
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.				
Specificity/Sensitivity		VAMP1 Antibody recognizes endogenous levels of total VAMP1 protein. This antibody does not cross-react with VAMP2.				
Species predicted to react based on 100% sequence homology		Monkey				
Source / Purification		Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues near the amino terminus of human VAMP1 protein. Antibodies are purified by protein A and peptide affinity chromatography.				
Background		Vesicle-associated membrane protein 1 (VAMP1), also called synaptobrevin 1, is part of the R-soluble N-ethylmaleimide-sensitive factor attachment protein receptor (SNARE) complex (1). The SNARE complex is involved in calcium regulated vesicular transport and membrane fusion (2). While related protein VAMP2 exhibits a wider distribution and is more abundant in the brain, VAMP1 is the main isoform in specific brain regions including the subthalamus nucleus zona incerta (1), the ostral periolivary region, and the retina (3). In addition, VAMP1 is involved in neurotransmitter release at the neuromuscular junction (4) and in the release of bioactive peptides from cardiac myocytes (5).				
Background References		 Raptis, A. et al. (2005) J Chem Neuroanat 30, 201-11. Sudhof, T.C. (2004) Annu Rev Neurosci 27, 509-47. Nystuen, A.M. et al. (2007) Neurogenetics 8, 1-10. Liu, Y. et al. (2011) J Physiol 589, 1603-18. Ferlito, M. et al. (2010) J Mol Cell Cardiol 49, 791-800. 				
Species Reactivity		Species reactivity is de	etermined by testin	g in at least one approve	ed application (e.g.,	western blot).
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X				

TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key W: Western Blotting IP: Immunoprecipitation

Cross-Reactivity Key H: Human M: Mouse R: Rat

Trademarks and Patents Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for

more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.