

Store at
-20C
#13256**CtBP2 Antibody**

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP	H M R Mk	Endogenous	47	Rabbit	#P56545	1488

Product Usage Information**Application**

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

CtBP2 Antibody recognizes endogenous levels of total CtBP2 protein. This antibody does not cross-react with the CtBP1 protein.

Species predicted to react based on 100% sequence homology

Bovine, Dog, Guinea Pig

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human CtBP2 protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

CtBP2 (carboxy-terminal binding protein-2) and its homolog CtBP1 are transcriptional co-repressors originally identified as proteins that bind the carboxy-terminus of the human adenovirus E1A protein (1-3). CtBP proteins are thought to play important roles in regulating various developmental pathways because deletion of CtBP2 leads to embryonic lethality at E10.5 and is correlated with axial patterning defects (4). CtBP proteins regulate various oncogenic signaling pathways as promoters of epithelial-mesenchymal transition, apoptosis antagonists, and tumor suppressor genes repressors (1,5). The CtBP protein transcription co-repression activity results from interactions with numerous transcription factors and chromatin modulators, including the polycomb group proteins (1,6,7). Depending on the context, CtBP proteins interact with a short amino acid sequence motif (PXDLS) to mediate repression of target genes through both histone deacetylase-dependent and independent mechanisms (6,8,9). CtBP proteins display a high sequence homology to the bacterial D-isomer-specific 2-hydroxyacid dehydrogenase enzymes. Research studies indicate that nuclear NADH levels regulate CtBP transcription repression activities, as NADH binding is required for CtBP2 homodimerization and transcription co-repressor activity (6,9-11).

Background References

- Chinnadurai, G. (2009) *Cancer Res* 69, 731-4.
- Boyd, J.M. et al. (1993) *EMBO J* 12, 469-78.
- Katsanis, N. and Fisher, E.M. (1998) *Genomics* 47, 294-9.
- Hildebrand, J.D. and Soriano, P. (2002) *Mol Cell Biol* 22, 5296-307.
- Battaglia, S. et al. (2010) *Int J Cancer* 126, 2511-9.
- Chinnadurai, G. (2002) *Mol Cell* 9, 213-24.
- Sewalt, R.G. et al. (1999) *Mol Cell Biol* 19, 777-87.
- Molloy, D.P. et al. (1998) *J Biol Chem* 273, 20867-76.
- Schaeper, U. et al. (1995) *Proc Natl Acad Sci U S A* 92, 10467-71.
- Kumar, V. et al. (2002) *Mol Cell* 10, 857-69.
- Thio, S.S. et al. (2004) *Nucleic Acids Res* 32, 1836-47.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.