

TRIB2 (D8P2X) Rabbit mAb

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP	Reactivity: H M	Sensitivity: Endogenous	MW (kDa): 42	Source/Isotype: Rabbit IgG	UniProt ID: #Q92519	Entrez-Gene Id: 28951
-------------------------------	---------------------------	-----------------------------------	------------------------	--------------------------------------	-------------------------------	---------------------------------

Product Usage Information**Application**

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:200

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

TRIB2 (D8P2X) Rabbit mAb recognizes endogenous levels of total TRIB2 protein. This antibody does not cross-react with other TRIBBLES family proteins.

Source / Purification

Monoclonal antibody is produced by immunizing animals with recombinant protein corresponding to full-length human TRIB2 protein.

Background

TRIBBLES proteins belong to a small family of serine-threonine kinase-like proteins characterized by the presence of a variant protein kinase motif (lacking a canonical ATP binding site), a MEK1 binding site, and a C-terminal COP1 site that binds ubiquitin ligase. The *tribbles* gene was first identified and characterized in *Drosophila* genetic screens for genes that regulate cell division, gastrulation, and oogenesis (1-3). Research studies in *Drosophila* suggested that TRIBBLES functions to coordinate cell division by regulating turnover of the cell cycle protein String/cdc25. In contrast to the *Drosophila* genome, which contains a single *tribbles* gene, the genomes of mice and humans encode three known TRIBBLES proteins (TRIB1-3), which exhibit both distinct and overlapping patterns of expression and functions (4). For example, TRIB1 and TRIB2, but not TRIB3, were reported to promote degradation of the basic region-leucine zipper transcription factor C/EBP α , a function that appears to be conserved from flies to humans (5,6). TRIB2 is overexpressed in a subset of human AML patient samples, downregulated in leukemic cells undergoing proliferation arrest (7), and positively regulated by the NOTCH signaling pathway in T cells (8), while retroviral-mediated overexpression of TRIB2 in mice was shown to induce transplantable leukemia (7). These findings collectively suggest that TRIB2 functions as an oncogene in the mammalian hematopoietic system (9).

Background References

- Grosshans, J. and Wieschaus, E. (2000) *Cell* 101, 523-31.
- Seher, T.C. and Leptin, M. (2000) *Curr Biol* 10, 623-9.
- Mata, J. et al. (2000) *Cell* 101, 511-22.
- Dobens, L.L. and Bouyain, S. (2012) *Dev Dyn* 241, 1239-48.
- Dedhia, P.H. et al. (2010) *Blood* 116, 1321-8.
- Rørth, P. et al. (2000) *Mol Cell* 6, 23-30.
- Keeshan, K. et al. (2006) *Cancer Cell* 10, 401-11.
- Hannon, M.M. et al. (2012) *Br J Haematol* 158, 626-34.
- Liang, K.L. et al. (2013) *Blood* 121, 4265-70.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: Human **M:** Mouse

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.