

Store at
-20C
#13701**SUZ12 (D39F6) XP[®] Rabbit mAb
(Biotinylated)**
Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W	H M R Mk	Endogenous	83	Rabbit IgG	#Q15022	23512

Product Usage Information**Application**

Western Blotting

Dilution

1:1000

StorageSupplied in 136 mM NaCl, 2.6 mM KCl, 12 mM sodium phosphate (pH 7.4) dibasic, 2 mg/ml BSA, and 50% glycerol. Store at -20°C. *Do not aliquot the antibodies.***Specificity/Sensitivity**SUZ12 (D39F6) XP[®] Rabbit mAb (Biotinylated) recognizes endogenous levels of total SUZ12 protein.**Species predicted to react based on 100% sequence homology**

Pig, Horse

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to the sequence of the human SUZ12 protein.

DescriptionThis Cell Signaling Technology antibody is conjugated to biotin under optimal conditions. The biotinylated antibody is expected to exhibit the same species cross-reactivity as the unconjugated SUZ12 (D39F6) XP[®] Rabbit mAb #3737.**Background**

The polycomb group (PcG) proteins contribute to the maintenance of cell identity, stem cell self-renewal, cell cycle regulation and oncogenesis by maintaining the silenced state of genes that promote cell lineage specification, cell death and cell-cycle arrest (1-4). PcG proteins exist in two complexes that cooperate to maintain long-term gene silencing through epigenetic chromatin modifications. The first complex, EED-EZH2, is recruited to genes by DNA-binding transcription factors and methylates histone H3 on Lys27. Methylation of Lys27 facilitates the recruitment of the second complex, PRC1, which ubiquitinylates histone H2A on Lys119 (5). Suppressor of Zeste 12 (SUZ12) is a component of the PRC2 complex, which together with Ezh2 and Eed is absolutely required for histone methyl-transferase activity (6). SUZ12 contains a C2H2 zinc finger domain similar to the zinc finger domains found in sequence-specific DNA binding proteins and may mediate the interaction between EZH2 and nucleosomes (6). SUZ12 is overexpressed in several human tumors, including tumors of the colon, breast and liver (7,8).

Background References

1. Boyer, L.A. et al. (2006) *Nature* 441, 349-53.
2. Lee, T.I. et al. (2006) *Cell* 125, 301-13.
3. Cao, R. et al. (2002) *Science* 298, 1039-43.
4. Müller, J. et al. (2002) *Cell* 111, 197-208.
5. Wang, H. et al. (2004) *Nature* 431, 873-8.
6. Cao, R. and Zhang, Y. (2004) *Mol Cell* 15, 57-67.
7. Kirmizis, A. et al. (2003) *Mol Cancer Ther* 2, 113-21.
8. Kirmizis, A. et al. (2004) *Genes Dev* 18, 1592-605.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.

Applications Key**W:** Western Blotting**Cross-Reactivity Key****H:** Human **M:** Mouse **R:** Rat **Mk:** Monkey**Trademarks and Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.