

Store at
-20C
#13705**U2AF1 (D6S3Q) Rabbit mAb**

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W	H M R Mk	Endogenous	35	Rabbit IgG	#Q01081	7307

Product Usage Information**Application**

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. *Do not aliquot the antibody.*

Specificity/Sensitivity

U2AF1 (D6S3Q) Rabbit mAb recognizes endogenous levels of total U2AF1 protein.

Species predicted to react based on 100% sequence homology

Hamster, Zebrafish, Bovine

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ala47 of human U2AF1 protein.

Background

U2 small nuclear RNA auxiliary factor 1 (U2AF1) is the small (35 kDa) subunit of the U2 auxiliary factor (U2AF) that plays an essential role in the splicing of pre-mRNA to generate functional mRNA transcripts. U2AF1 forms a heterodimer with the large (65 kDa) U2AF2 subunit to create the U2 auxiliary factor that recognizes the 3' splice site and facilitates spliceosome assembly (1-3). Research studies indicate that U2AF1 binds to the 3'-splice site consensus AG dinucleotide at the intron-exon boundary while U2AF2 recognizes and binds the polyprimidine tract upstream of the 3' splice site. These two steps ensure accurate spliceosome assembly at splice sites (4-6). Mutations in the corresponding *U2AF1* gene are associated with a type of hematopoietic stem cell disorder known as myelodysplastic syndrome (MDS), which can be characterized by low blood counts, anemia, and enhanced acute myeloid leukemia risk (7-9). Somatic U2AF1 mutations frequently affect highly conserved zinc finger protein regions that result in defective pre-mRNA splicing of genes involved in cell cycle progression and RNA processing pathways, contributing to MDS pathogenesis (7,10).

Background References

1. Jurica, M.S. and Moore, M.J. (2003) *Mol Cell* 12, 5-14.
2. Kielkopf, C.L. et al. (2001) *Cell* 106, 595-605.
3. Zamore, P.D. and Green, M.R. (1989) *Proc Natl Acad Sci U S A* 86, 9243-7.
4. Merendino, L. et al. (1999) *Nature* 402, 838-41.
5. Wu, S. et al. (1999) *Nature* 402, 832-5.
6. Zamore, P.D. et al. (1992) *Nature* 355, 609-14.
7. Graubert, T.A. et al. (2012) *Nat Genet* 44, 53-7.
8. Hahn, C.N. and Scott, H.S. (2012) *Nat Genet* 44, 9-10.
9. Yoshida, K. et al. (2011) *Nature* 478, 64-9.
10. Przychodzen, B. et al. (2013) *Blood* 122, 999-1006.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's

terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com
For Research Use Only. Not for Use in Diagnostic Procedures.