

CAMLG (D5L9J) Rabbit mAb

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP	H M R Hm Mk	Endogenous	32	Rabbit IgG	#P49069	819

Product Usage Information**Application**

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:200

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

CAMLG (D5L9J) Rabbit mAb recognizes endogenous levels of total CAMLG protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Leu110 of human CAMLG protein.

Background

The calcium signal-modulating cyclophilin ligand (CAMLG) is a multi-pass endoplasmic reticulum (ER) membrane protein that plays an important role in calcium-mediated signal transduction. The CAMLG protein was first identified in T lymphocytes as a cyclophilin B-binding protein that regulates calcium influx following T-cell receptor (TCR) activation (1). Research studies indicate that CAMLG overexpression activates the transcription factors NFAT and NF-IL2A and leads to increased IL-2 gene transcription, providing additional evidence that CAMLG plays an important role in TCR signal transduction. CAMLG negatively regulates the non-receptor protein kinase Lck, which is critical for the thymocyte selection process (2). Thymocytes deficient for CAMLG fail to undergo normal development, accumulate reactive oxygen species, and exhibit increased apoptosis in response to cytotoxic stimuli (3). The CAMLG protein also acts as a receptor of TRC40, an ATPase that targets newly synthesized proteins in the secretory pathway to the ER membrane. CAMLG interacts with the protein insertion receptor WRB to form the TRC40 receptor complex that is responsible for insertion of tail-anchored proteins into the ER membrane (4).

Background References

1. Bram, R.J. and Crabtree, G.R. (1994) *Nature* 371, 355-8.
2. Tran, D.D. et al. (2005) *Immunity* 23, 139-52.
3. Edgar, C.E. et al. (2010) *Cell Death Differ* 17, 1566-76.
4. Yamamoto, Y. and Sakisaka, T. (2012) *Mol Cell* 48, 387-97.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat **Hm:** Hamster **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any

purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.