

#1406

Store at -20°C

Flt3 (Tyr919) Biotinylated Peptide

1.25 ml at 6 µM



Cell Signaling
TECHNOLOGY®

Orders ■ 877-616-CELL (2355)
orders@cellsignal.com

Support ■ 877-678-TECH (8324)
info@cellsignal.com

Web ■ www.cellsignal.com

new 09/06

This product is for *in vitro* research use only and is not intended for use in humans or animals.

Description: This biotinylated peptide contains the residues surrounding Tyr919 of Flt3. It was generated for use in CST's HTScan® kinase assay kits, but may also serve as a substrate in other heterogeneous or homogeneous kinase assays.

Peptide Core Sequence: EEIY*IIM

Molecular Weight: 1779 daltons

Quality Control: The quality of the biotinylated peptide was evaluated by reverse-phase HPLC and by mass spectrometry.

Directions for Use: The phosphorylated form of the peptide can be detected with Phospho-Tyrosine mAb (P-Tyr-100) #9411. Sample kinase assay protocol is attached.

Storage: Supplied in 0.0001% DMSO. Store at -20°C.

Companion Products:

Tyrosine Kinase Substrate Screening Kit #7450

Phospho-Tyrosine Mouse mAb (P-Tyr-100) #9411

TrkA Kinase #7422

Protocol for Tyrosine Kinase Assay

IMPORTANT: Use of an automated microplate washer as well as centrifugation of plates when appropriate, greatly improves reproducibility.

A Additional Solutions and Reagents (Not included)

1. **Wash Buffer:** 1X PBS, 0.05% Tween-20 (PBS/T)
2. Bovine Serum Albumin (BSA)
3. **Stop Buffer:** 50 mM EDTA pH 8
4. Phospho-Tyrosine mAb (P-Tyr-100) #9411
5. Kinase Buffer (4X) #9805
6. ATP (10 mM) #9804
7. DTT (1.25M)
8. Kinase (See companion products)
9. DELFIA® Europium-labeled Anti-mouse IgG (PerkinElmer Life Sciences #AD0124)
10. DELFIA® Enhancement Solution (PerkinElmer Life Sciences #1244-105)
11. DELFIA® Streptavidin coated, 96-well, yellow plate (PerkinElmer Life Sciences AAAND-0005)

DELFIA® is a registered trademark of PerkinElmer Life Sciences

B Suggested Protocol for 100 Assays

1. Add 100 µl 10 mM ATP to 1.25 ml 6-12 µM substrate peptide. Adjust the mixture with dH₂O to 2.5 ml to make 2X ATP/substrate cocktail ([ATP]=400 µM, [substrate] = 3-6 µM).
2. Transfer enzyme from -80°C to ice. Allow enzyme to thaw on ice.
3. **Microcentrifuge briefly at 4°C to bring liquid to the bottom of the vial. Return immediately to ice.**
4. Add 10 ml of DTT (1.25M) to 2.5 ml of 4X HTScan® tyrosine kinase buffer (240 mM HEPES pH 7.5, 20 mM MgCl₂, 20 mM MnCl₂, 12 mM Na₃VO₄) to make 4xDTT/Kinase buffer.
5. Dilute enzyme in 1.25 ml of 4X DTT/Kinase buffer to make 4X reaction cocktail (enzyme)=0.8-8.0 ng/µl in 4X DTT/Kinase buffer).
6. Add 12.5 µl of the 4X reaction cocktail to 12.5 µl/well of prediluted compound of interest (usually around 10 µM) and incubate for 5 minutes at room temperature.
7. Add 25 µl of 2X ATP/substrate cocktail to 25 µl/well preincubated reaction cocktail/compound.

Final Assay Conditions for a 50 µl Reaction

- 60 mM HEPES (pH7.5)
 - 5 mM MgCl₂
 - 5 mM MnCl₂
 - 3 mM Na₃VO₄
 - 1.25 mM DTT
 - 200 µM ATP
 - 1.5-3 µM peptide
 - 10-100 ng kinase
8. Incubate reaction plate at room temperature for 30 minutes.
 9. Add 50 µl/well Stop Buffer (50 mM EDTA, pH 8) to stop the reaction.
 10. Transfer 25 µl of each reaction to a 96-well streptavidin-coated plate containing 75 µl dH₂O/well and incubate at room temperature for 60 minutes.
 11. Wash three times with 200 µl/well PBS/T.
 12. Dilute primary antibody (Phospho-Tyrosine mAb (P-Tyr-100) #9411) in PBS/T with 1% BSA. *Add 100 µl/well primary antibody.
 13. Incubate at 37°C for 120 minutes.
 14. Wash three times with 200 µl/well PBS/T.
 15. Dilute Europium labeled secondary antibody in PBS/T with 1% BSA. **Add 100 µl/well diluted antibody.
 16. Incubate at room temperature for 30 minutes.
 17. Wash five times with 200 µl/well PBS/T.
 18. Add 100 µl/well DELFIA® Enhancement Solution.
 19. Incubate at room temperature for 5 minutes.
 20. Detect 615 nm fluorescence emission with appropriate Time-Resolved Plate Reader.

*Recommended antibody dilution factor:

Primary antibody:
 Mouse mAb: 1:500
 Rabbit mAb or poly-clone antibody: 1:1000

**Secondary antibody:

DELFIA® Europium-labeled Anti-mouse IgG: 1:500
 DELFIA® Europium-labeled Anti-rabbit antibody: 1:1000