

Store at  
-20°C  
#14066

# SUMO-2/3 (18H8) Rabbit mAb (HRP Conjugate)



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W	H M R	Endogenous	Rabbit IgG	#P55854, #P61956	6612, 6613

## Product Usage Information

### Application

Western Blotting

### Dilution

1:1000

## Storage

Supplied in 136 mM NaCl, 2.6 mM KCl, 12 mM sodium phosphate (pH 7.4) dibasic, 2 mg/ml BSA, and 50% glycerol. Store at -20°C. Do not aliquot the antibodies.

## Specificity/Sensitivity

SUMO-2/3 (18H8) Rabbit mAb (HRP Conjugate) detects endogenous levels of SUMO-2/3 protein. It does not cross-react with recombinant SUMO-1 protein.

## Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide from the amino terminus of human SUMO-3 protein.

## Description

This Cell Signaling Technology antibody is conjugated to the carbohydrate groups of horseradish peroxidase (HRP) via its amine groups. The HRP conjugated antibody is expected to exhibit the same species cross-reactivity as the unconjugated SUMO-2/3 (18H8) Rabbit mAb #4971.

## Background

Small ubiquitin-related modifier 1, 2 and 3 (SUMO-1, -2 and -3) are members of the ubiquitin-like protein family (1). The covalent attachment of the SUMO-1, -2 or -3 (SUMOylation) to target proteins is analogous to ubiquitination. This post-translational modification is a reversible, multi-step process that is initiated by cleaving a precursor protein to a mature protein. Mature SUMO-1, -2 or -3 is then linked to the activating enzyme E1, conjugated to E2 and in conjunction with E3, SUMO-1, -2 or -3 is ligated to the target protein (2). Ubiquitin and the individual SUMO family members are all targeted to different proteins with diverse biological functions. Ubiquitin predominantly regulates degradation of its target (1). In contrast, SUMO-1 is conjugated to RanGAP, PML, p53 and IκB-α to regulate nuclear trafficking, formation of subnuclear structures, regulation of transcriptional activity and protein stability (3-7). SUMO-2/-3 forms poly-(SUMO) chains, is conjugated to topoisomerase II and APP, regulates chromosomal segregation and cellular responses to environmental stress, and plays a role in the progression of Alzheimer disease (8-11).

## Background References

- Schwartz, D.C. and Hochstrasser, M. (2003) *Trends Biochem. Sci.* 28, 321-8.
- Kim, K.I. et al. (2002) *J. Cell Physiol.* 191, 257-68.
- Matunis, M.J. et al. (1996) *J. Cell Biol.* 135, 1457-70.
- Duprez, E. et al. (1999) *J. Cell Sci.* 112, 381-93.
- Gostissa, M. et al. (1999) *EMBO J.* 18, 6462-74.
- Rodriguez, M.S. et al. (1999) *EMBO J.* 18, 6455-61.
- Desterro, J.M. et al. (1998) *Mol. Cell* 2, 233-9.
- Tatham, M.H. et al. (2001) *J. Biol. Chem.* 276, 35368-74.
- Azuma, Y. et al. (2003) *J. Cell Biol.* 163, 477-87.
- Li, Y. et al. (2003) *Proc. Natl. Acad. Sci. USA* 100, 259-64.
- Saitoh, H. and Hinchey, J. (2000) *J. Biol. Chem.* 275, 6252-8.

## Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

## Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

## Applications Key

**W:** Western Blotting

## Cross-Reactivity Key

**H:** Human **M:** Mouse **R:** Rat

## Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

## Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.