

**p70 S6 Kinase 2 Antibody**

**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP	H M R	Endogenous	60	Rabbit	#Q9UBS0	6199

**Product Usage Information****Application**

Western Blotting  
Immunoprecipitation

**Dilution**

1:1000  
1:50

**Storage**

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

**Specificity/Sensitivity**

p70 S6 Kinase 2 Antibody recognizes endogenous levels of total p70 S6 kinase 2 (S6K2) protein. This antibody does not cross-react with p70 S6 kinase 1 (S6K1) protein.

**Source / Purification**

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Pro435 of human p70 S6 kinase 2 protein. Antibodies are purified by protein A and peptide affinity chromatography.

**Background**

p70 S6 kinase is a mitogen activated Ser/Thr protein kinase that is required for cell growth and G1 cycle progression (1,2). p70 S6 kinase phosphorylates the S6 protein of the 40S ribosomal subunit and is involved in translational control of 5' oligopyrimidine tract mRNAs (1). A second isoform, p85 S6 kinase, is derived from the same gene and is identical to p70 S6 kinase except for 23 extra residues at the amino terminus, which encode a nuclear localizing signal (1). Both isoforms lie on a mitogen activated signaling pathway downstream of phosphoinositide-3 kinase (PI-3K) and the target of rapamycin, FRAP/mTOR, a pathway distinct from the Ras/MAP kinase cascade (1). The activity of p70 S6 kinase is controlled by multiple phosphorylation events located within the catalytic, linker and pseudosubstrate domains (1). Phosphorylation of Thr229 in the catalytic domain and Thr389 in the linker domain are most critical for kinase function (1). Phosphorylation of Thr389, however, most closely correlates with p70 kinase activity *in vivo* (3). Prior phosphorylation of Thr389 is required for the action of phosphoinositide 3-dependent protein kinase 1 (PDK1) on Thr229 (4,5). Phosphorylation of this site is stimulated by growth factors such as insulin, EGF and FGF, as well as by serum and some G-protein-coupled receptor ligands, and is blocked by wortmannin, LY294002 (PI-3K inhibitor) and rapamycin (FRAP/mTOR inhibitor) (1,6,7). Ser411, Thr421 and Ser424 lie within a Ser-Pro-rich region located in the pseudosubstrate region (1). Phosphorylation at these sites is thought to activate p70 S6 kinase via relief of pseudosubstrate suppression (1,2). Another LY294002 and rapamycin sensitive phosphorylation site, Ser371, is an *in vitro* substrate for mTOR and correlates well with the activity of a partially rapamycin resistant mutant p70 S6 kinase (8).

p70 S6 kinase 2 (S6K2) exhibits high homology in the kinase domain and adjacent regulatory region with p70 S6 kinase (S6K1). Similar to S6K1, S6K2 displays both mitogen-dependent and rapamycin-sensitive S6 kinase activity (9,10). S6K2 has been shown to have redundant as well as distinct functions from S6K1 (11,12). Research studies show that S6K2 is commonly expressed at higher levels in tumor samples than in corresponding normal tissues, and may promote cancer cell survival via AKT activation (13,14).

**Background References**

- Pullen, N. and Thomas, G. (1997) *FEBS Lett* 410, 78-82.
- Dufner, A. and Thomas, G. (1999) *Exp Cell Res* 253, 100-9.
- Weng, Q.P. et al. (1998) *J Biol Chem* 273, 16621-9.
- Pullen, N. et al. (1998) *Science* 279, 707-10.
- Alessi, D.R. et al. (1998) *Curr Biol* 8, 69-81.
- Polakiewicz, R.D. et al. (1998) *J Biol Chem* 273, 23534-41.
- Fingar, D.C. et al. (2002) *Genes Dev* 16, 1472-87.
- Saitoh, M. et al. (2002) *J Biol Chem* 277, 20104-12.
- Gout, I. et al. (1998) *J Biol Chem* 273, 30061-4.
- Lee-Fruman, K.K. et al. (1999) *Oncogene* 18, 5108-14.
- Pende, M. et al. (2004) *Mol Cell Biol* 24, 3112-24.
- Pardo, O.E. et al. (2006) *EMBO J* 25, 3078-88.
- Sridharan, S. and Basu, A. (2011) *Cancer Res* 71, 2590-9.
- Pardo, O.E. and Seckl, M.J. (2013) *Front Oncol* 3, 191.

<b>Species Reactivity</b>	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
<b>Western Blot Buffer</b>	<b>IMPORTANT:</b> For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.
<b>Applications Key</b>	<b>W:</b> Western Blotting <b>IP:</b> Immunoprecipitation
<b>Cross-Reactivity Key</b>	<b>H:</b> Human <b>M:</b> Mouse <b>R:</b> Rat
<b>Trademarks and Patents</b>	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit <a href="http://cellsignal.com/trademarks">cellsignal.com/trademarks</a> for more information.
<b>Limited Uses</b>	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.  Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.