

Store at
-20°C

#14163

HMGB2 (D1P9V) Rabbit mAb

www.cellsignal.com

Support: 877-678-TECH (8324)
info@cellsignal.comOrders: 877-616-CELL (2355)
orders@cellsignal.comEntrez-Gene ID #3148
UniProt ID #P26583

rev. 07/07/14


For Research Use Only. Not For Use In Diagnostic Procedures.

Applications W Endogenous	Species Cross-Reactivity* H, M, R, Mk, (Hm, B, Dg, Hr, guinea pig)	Molecular Wt. 28 kDa	Isotype Rabbit IgG**
---------------------------------	--	-------------------------	-------------------------

Background: High mobility group protein B2 (HMGB2) belongs to a family of highly conserved proteins that contain HMG box domains (1,2). All three family members (HMGB1, HMGB2, and HMGB3) contain two HMG box domains and a C-terminal acidic domain. HMGB1 is a widely expressed and highly abundant protein (2). HMGB2 is widely expressed during embryonic development, but it is restricted to lymphoid organs and testis in adult animals (3). HMGB3 is only expressed during embryogenesis (4). While expression varies, the biochemical properties of the different family members may be indistinguishable. The HMG box domains facilitate the binding of HMGB proteins to the minor groove of DNA, which results in local bending of the DNA double helix (1,2). HMGB proteins are recruited by and help facilitate the assembly of site-specific DNA binding proteins to their cognate binding sites in chromatin. For example, HMGB1 and HMGB2 facilitate the binding of Hox proteins, Oct proteins, p53, Rel proteins, and steroid hormone receptor proteins to their target gene promoters (1,2). Furthermore, HMGB2 interacts with RAG1 to facilitate RAG complex binding to the recombinant signal sequence (RSS) and stimulate DNA-bending and subsequent VDJ cleavage at antigen receptor genes (5,6). In addition to their functions in the nucleus, HMGB proteins play a significant role in extracellular signaling associated with inflammation. HMGB2 is secreted by myeloid cells and promotes proliferation and migration of endothelial cells by binding to the receptor for advanced glycation endproducts (RAGE) (7). Research studies have shown that HMGB2 overexpression in hepatocellular carcinoma is associated with poor prognosis and shorter survival time (8).

Specificity/Sensitivity: HMGB2 (D1P9V) Rabbit mAb recognizes endogenous levels of total HMGB2 protein. This antibody does not cross-react with other HMGB proteins, including HMGB1 and HMGB3.

Source/Purification: Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Glu169 of human HMGB2 protein.



Western blot analysis of extracts from various cell lines using HMGB2 (D1P9V) Rabbit mAb.

Storage: Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. *Do not aliquot the antibody.*

***Species cross-reactivity is determined by western blot.**

****Anti-rabbit secondary antibodies must be used to detect this antibody.**

Recommended Antibody Dilutions:

Western blotting 1:1000

For product specific protocols please see the web page for this product at www.cellsignal.com.

Please visit www.cellsignal.com for a complete listing of recommended complementary products.

Background References:

- (1) Thomas, J.O. and Travers, A.A. (2001) *Trends Biochem Sci* 26, 167-74.
- (2) Müller, S. et al. (2004) *J Intern Med* 255, 332-43.
- (3) Ronfani, L. et al. (2001) *Development* 128, 1265-73.
- (4) Vaccari, T. et al. (1998) *Genomics* 49, 247-52.
- (5) Zhang, M. and Swanson, P.C. (2009) *BMC Mol Biol* 10, 24.
- (6) Aidinis, V. et al. (1999) *Mol Cell Biol* 19, 6532-42.
- (7) Pusterla, T. et al. (2009) *Autoimmunity* 42, 308-10.
- (8) Kwon, J.H. et al. (2010) *Clin Cancer Res* 16, 5511-21.

Tween® is a registered trademark of ICI Americas, Inc.

IMPORTANT: For western blots, incubate membrane with diluted antibody in 5% w/v BSA, 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

© 2014 Cell Signaling Technology, Inc.
Cell Signaling Technology® is a trademark of Cell Signaling Technology, Inc.

Thank you for your recent purchase. If you would like to provide a review visit www.cellsignal.com/comments.



Applications: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide Species Cross-Reactivity: H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—Horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.